

**One Earth, Volume 6**

**Supplemental information**

**Many risky feedback loops**

**amplify the need for climate action**

**William J. Ripple, Christopher Wolf, Timothy M. Lenton, Jillian W. Gregg, Susan M. Natali, Philip B. Duffy, Johan Rockström, and Hans Joachim Schellnhuber**

**Supplemental Tables**

**Table S1. Summary of 41 climate feedback loops.** The first column indicates the primary system or process involved in the loop (e.g., clouds). “Effect of climate change” describes how climate change alters the system (an active process) and “Effect on climate change” describes how the response of the system contributes to climate change. Subsequent columns indicate overall/net feedback direction (positive loops shown are shown in red), strength, and type respectively. References for strength estimates are listed in the first column (“Feedback name”) or in the footnotes (when the feedback has multiple references). Type refers only to the dominant type or types associated with the feedback loop. When the overall feedback direction is unknown, this is indicated with a “?” symbol. Loops are divided into two categories: physical (no significant biological or human component [aside from anthropogenic climate change]) and biological (includes a biological component but no human component). Feedback loops that involve potential tipping elements identified in Table 1 of Lenton et al.<sup>1</sup> are marked with asterisks (\*) and feedback loops that are covered in Fig. TS.17 (feedbacks overview) or 5.29 (biogeochemical feedbacks) of IPCC<sup>2</sup> with daggers (†). Some feedback loops may be highly uncertain or speculative; especially, those for which feedback strength estimates (or confidence intervals, etc.) were unavailable. Thus, it should not be assumed that all loops in the table are equally important in any sense. Feedback impacts occur on various time scales ranging from short (e.g., months or years) to very long (e.g., millennia); feedbacks we believe to be exceptionally slow (i.e., millennia) are marked with double daggers (‡). There are some instances of overlap among feedback loops (e.g., “soil carbon (other)” and “permafrost”), which are identified with footnotes. In these cases, strength estimates are not necessarily additive (see Supplemental Experimental Procedures).

<b>Feedback name</b>	<b>Effect of climate change</b>	<b>Effect on climate change</b>	<b>+/-</b>	<b>Approximate strength</b>	<b>Type</b>
<b>20 physical (abiotic) feedback loops</b>					
<b>1. Planck (black body radiation)<sup>† 3,4</sup></b>	Increasing mean surface and atmosphere temperature	Increasing heat loss due to emitted radiation	-	-3.22 (-3.4 to -3.0) W/m <sup>2</sup> /K <sup>a</sup>	Other
<b>2. Water vapor<sup>† 5,6,3</sup></b>	Increasing water vapor content due to warming	Increasing greenhouse effect (water vapor is a greenhouse gas)	+	2.04 W/m <sup>2</sup> /K <sup>b</sup>	Water vapor
<b>3. Sea ice albedo<sup>*† 8-14 c</sup></b>	Sea ice melting <sup>d</sup> or not forming	Decreasing albedo	+ <sup>e</sup>	0.31 W/m <sup>2</sup> /K <sup>f</sup>	Surface albedo <sup>g</sup>
<b>4. Glacier and ice sheet albedo<sup>*†‡ 21,22</sup></b>	Increasing glacier & ice sheet melting and marine ice sheet instability <sup>23-25</sup>	Decreasing albedo	+		Surface albedo
<b>5. Sea level rise<sup>† 26,27</sup></b>	Rising sea levels (caused by ice melting)	Decreasing albedo due to coastal regions being submerged	+		Surface albedo
<b>6. Snow cover<sup>† 28-30</sup></b>	Snow metamorphosis and decreasing snow cover	Decreasing albedo	+	0.03 to 0.16 W/m <sup>2</sup> /K <sup>h</sup>	Surface albedo

Feedback name	Effect of climate change	Effect on climate change	+/-	Approximate strength	Type
<b>7. Clouds</b> <sup>†</sup> 8,31–33,20,3,4	Changing cloud distribution (extent and height) and optical properties	Changing cloud albedo and greenhouse effect	+	0.45 W/m <sup>2</sup> /K (σ = 0.33 W/m <sup>2</sup> /K) <sup>i</sup>	Cloud
<b>8. Dust</b> <sup>†</sup> 34–37	Changing dust aerosol abundance <sup>j</sup>	Changing planetary albedo and greenhouse effect	?	-0.004 ± 0.007 W/m <sup>2</sup> /K <sup>k</sup>	Aerosol
<b>9. Other aerosols</b> <sup>†</sup> 34,38,3,39,36,37	Changing atmospheric aerosol (e.g., sulfate, sea salt) concentrations	Changing planetary albedo and greenhouse effect	?	<sup>l</sup>	Aerosol
<b>10. Ocean stratification</b> <sup>40</sup> ,41 m	Ocean warming resulting in increasing stratification	Decreasing carbon uptake by ocean <sup>n</sup>	+		Ocean
<b>11. Ocean circulation</b> <sup>*</sup> 42–47 o	Slowing of the Atlantic Meridional Overturning Circulation (AMOC) <sup>p</sup>	Changing surface temperature	?	Independent short term strength not applicable <sup>q</sup>	Ocean
<b>12. Ocean solubility pump</b> <sup>49,50</sup>	Increasing atmospheric CO <sub>2</sub> levels	Decreasing absorption of CO <sub>2</sub> by ocean <sup>r</sup>	+	0.16 W/m <sup>2</sup> /K <sup>s</sup>	Ocean
<b>13. Methane hydrates</b> <sup>*†</sup> 52–57	Increasing methane hydrate dissociation rates	Increasing release of methane into the atmosphere <sup>t</sup>	+		Ocean
<b>14. Lapse rate</b> <sup>†</sup> 8,59,3,4	Changing relationships between temperature and altitude	Increasing heat loss due to emitted radiation	-	-0.68 to -0.23 W/m <sup>2</sup> /K <sup>u</sup>	Other
<b>15. Glacier and ice sheet elevation</b> <sup>†</sup> 23,60–62 v	Decreasing glacier and ice sheet mass balance and elevation	Increasing glacier and ice sheet melting (due to increasing surface temperatures), decreasing albedo	+		Other
<b>16. Antarctic rainfall</b> <sup>†</sup> 63	Decreasing Antarctic ice sheet extent leading to increasing precipitation	Decreasing albedo, increasing deep ocean warming	+		Other
<b>17. Sea ice growth</b> <sup>20,64 w</sup>	Decreasing sea ice thickness, increasing open-water fraction, decreasing insulation	Increasing growth rate of thin ice	-		Other
<b>18. Ozone</b> <sup>66,67,20,3,37 x</sup>	Strengthening of the Brewer-Dobson circulation, increasing stratosphere-to-troposphere transport	Decreasing lower tropical stratospheric ozone, increasing middle and extratropical stratospheric ozone, changing tropospheric ozone concentration	?	-0.064 (-0.08 to 0.04) W/m <sup>2</sup> /K <sup>y</sup>	Other

Feedback name	Effect of climate change	Effect on climate change	+/-	Approximate strength	Type
<b>19.</b> Atmospheric chemical reaction rates <sup>†20,36,37,68z</sup>	Changing chemical reaction rates in the atmosphere <sup>aa</sup>	Changing greenhouse effect	?	<sup>bb</sup>	Other
<b>20.</b> Chemical weathering <sup>‡70</sup>	Increasing carbonate and silicate weathering rates	Increasing CO <sub>2</sub> taken out of the atmosphere	-		Geological
<b>21 biological feedback loops</b>					
<b>21.</b> Peatlands <sup>71-73</sup>	Decreasing soil organic carbon due to lowering of water table, increasing vulnerability to fire, increasing metabolization	Increasing release of CO <sub>2</sub> into the atmosphere, decreasing carbon sequestration, changing methane emissions	+		Wetlands
<b>22.</b> Wetlands (expansion) <sup>†36,74,75</sup>	Increasing precipitation <sup>74</sup> and boreal near-surface soil moisture <sup>75</sup> potentially leading to expansion of wetlands <sup>cc</sup>	Increasing CO <sub>2</sub> sequestration (negative feedback) and methane emissions <sup>75-77</sup> (positive feedback)	+	0.16 ± 0.03 W/m <sup>2</sup> /K <sup>dd</sup>	Wetlands
<b>23.</b> Freshwater ecosystems <sup>†78,74,79-81 ee</sup>	Increasing aquatic plant growth rates (negative feedback) and microbial methane production (positive feedback)	Increasing methane emissions	+		Freshwater
<b>24.</b> Forest dieback <sup>*82,83,57,84</sup>	Dieback of Amazon, <sup>ff</sup> boreal, and other forests <sup>gg</sup>	Loss of sequestration, change in albedo, decreasing evapotranspiration	+		Trees
<b>25.</b> Northern greening <sup>93-99,20,100</sup>	Potential expansion of high latitude/elevation forests & woody vegetation into tundra, <sup>hh</sup> increasing Arctic/ northern vegetation <sup>ii</sup> (warmer & longer growing seasons, nutrient fertilization)	Increasing CO <sub>2</sub> sequestration, decreasing albedo <sup>jj</sup>	+		Trees/vegetation
<b>26.</b> Insect outbreaks (forests) <sup>103-105 kk</sup>	Changing insect distributions & abundances; decreased host tree defense	Loss of sequestration, change in albedo	+		Trees

Feedback name	Effect of climate change	Effect on climate change	+/-	Approximate strength	Type
<b>27. Wildfire</b> <sup>†</sup> 34,106–109 ll	Increasing fire frequency and/or severity and/or extent in some regions	Increasing CO <sub>2</sub> emissions, loss of sequestration, <sup>mm</sup> change in albedo	+	-0.10 to 0.44 W/m <sup>2</sup> by 2100 <sup>nn</sup>	Trees/aerosol
<b>28. Biogenic volatile organic compounds (BVOCs)</b> <sup>†</sup> 34,20,112,36,37	Changing BVOC emission rates	Decreasing greenhouse effect, increasing tropospheric ozone	-	-0.05 (-0.22 to 0.12) W/m <sup>2</sup> /K <sup>oo</sup>	Aerosol
<b>29. Soil carbon (other)</b> <sup>113–117 pp</sup>	Increasing loss of soil carbon to the atmosphere <sup>qq</sup>	Increasing CO <sub>2</sub> emissions	+		Soil
<b>30. Soil nitrous oxide</b> <sup>†</sup> 118–120,108 rr	Accelerated decomposition and changing soil microbial activity affecting substrate availability for denitrification	Increasing N <sub>2</sub> O emissions	+	1 Tg N/yr/K <sup>ss</sup>	Soil
<b>31. Permafrost</b> <sup>*†</sup> 123–130,57,131–133,108	Increasing thawing and decomposition <sup>tt</sup>	Increasing GHG (CO <sub>2</sub> and methane) emissions	+	0.03 to 0.29 W/m <sup>2</sup> /K <sup>uu</sup>	Soil
<b>32. Evapotranspiration from soils and plants</b> <sup>20</sup>	Increasing evaporation from soils, plants open stomata less widely <sup>vv</sup>	Decreasing latent heat flux (leading to warming)	+		Soil/vegetation
<b>33. Microbial respiration (other)</b> <sup>134,135 ww</sup>	Increasing respiration rates for many prokaryotic microbes	Increasing net CO <sub>2</sub> and CH <sub>4</sub> emissions due to microbes	+		Microbes
<b>34. Plant stress</b> <sup>136</sup>	Increasing chronic & extreme thermal and moisture stress	Increasing plant <sup>xx</sup> mortality leading to decreasing CO <sub>2</sub> sequestration	+		Vegetation
<b>35. Desertification</b> <sup>137,138 yy</sup>	Increasing chronic aridification & hotter drought stress extremes leading to expanding deserts	Decreasing CO <sub>2</sub> sequestration, increasing CO <sub>2</sub> emissions, <sup>zz</sup> and increasing albedo	+		Vegetation
<b>36. Sahara and Sahel greening</b> <sup>*</sup> 140,139 aaa	Possibly increasing rainfall in Sahara and Sahel <sup>bbb</sup>	Increasing CO <sub>2</sub> sequestration by vegetation, decreasing albedo	-		Vegetation
<b>37. CO<sub>2</sub> fertilization</b> 141,83,142,20,108	Increasing CO <sub>2</sub> concentration possibly leading to increasing net primary productivity (NPP) <sup>ccc</sup>	Possibly increasing carbon uptake by vegetation <sup>ddd</sup>	-	approx. 23% increase in forest NPP at 550 ppm CO <sub>2</sub> <sup>eee</sup>	Vegetation

Feedback name	Effect of climate change	Effect on climate change	+/-	Approximate strength	Type
<b>38. Coastal productivity</b> <sup>148-152</sup>	Increasing degradation of coastal ecosystems (e.g., mangroves, seagrass beds, salt marshes) due to heat stress and loss of coral reef protection	Decreasing carbon sequestration by coastal ecosystems	+		Coastal
<b>39. Ocean metabolic rates</b> <sup>57,153</sup>	Potentially increasing phytoplankton <sup>fff</sup> /bacterial respiration rates	Increasing CO <sub>2</sub> released into the atmosphere	+	approx. 0.02 °C by 2100 <sup>ggg</sup>	Ocean
<b>40. Ocean biological pump</b> <sup>155-158,108</sup>	Increasing CO <sub>2</sub> in ocean, ocean acidification, warming, decreasing upwelling	Changing effectiveness of ocean as a carbon sink, decreasing CO <sub>2</sub> uptake due to reduced primary productivity in upper layers of ocean <sup>hhh</sup>	? <sup>iii</sup>		Ocean
<b>41. Phytoplankton dimethyl sulfide (DMS)</b> <sup>† 160-163</sup>	Changing DMS emissions from plankton (due to ocean acidification, <sup>jjj</sup> increasing stratification, etc.)	Changing cloud longevity and albedo (leading to changing radiative forcing)	? <sup>kkk</sup>	0.005 (0.0 to 0.01) W/m <sup>2</sup> /K <sup>lll</sup>	Ocean/aerosol/albedo

<sup>a</sup> Central estimate and “very likely” interval from Table 7.10 of Forster et al.<sup>4</sup>

<sup>b</sup> Average feedback strength between 2003 and 2008 estimated by Dessler et al.<sup>5</sup> Note that there is a positive feedback associated with stratospheric water vapor in particular (approximate strength 0.3 W/m<sup>2</sup>/K according to Dessler et al.<sup>7</sup>).

<sup>c</sup> We have flagged this loop as being associated with a potential tipping point following Lenton et al.<sup>1</sup>, although evidence may be limited.<sup>9,15</sup>

<sup>d</sup> Both Arctic sea ice and Greenland ice sheet melting could potentially be accelerated by the jet stream slowing and becoming wavier due to climate change<sup>16-18</sup> – another possible feedback loop (but see Blackport and Screen<sup>19</sup>)

<sup>e</sup> However, this positive feedback may be countered by negative sea ice feedbacks – see “sea ice growth rate” feedback loop and Heinze et al.<sup>20</sup>

<sup>f</sup> Estimate for the entire globe obtained from Cao et al. 2015<sup>12</sup>

<sup>g</sup> Total surface albedo feedback strength estimated to be 0.30 W/m<sup>2</sup>/K ( $\sigma = 0.015$  W/m<sup>2</sup>/K) by Sherwood et al.<sup>3</sup>

<sup>h</sup> Strength estimate from Qu and Hall<sup>29</sup>

<sup>i</sup> Estimate from Table 1 of Sherwood et al.<sup>3</sup> While most models show a net positive feedback for clouds, some models exhibit a net negative cloud feedback. The net cloud feedback can be decomposed<sup>3</sup> – in short, increasing cloudiness means that high altitude clouds will trap more heat (positive feedback) while low altitude clouds will reflect more sunlight (negative feedback). For reference, the AR6 assessed range central estimate is 0.42 W/m<sup>2</sup>/K and the “very likely” interval is (-0.10 to 0.60) W/m<sup>2</sup>/K.<sup>4</sup>

<sup>j</sup> This feedback may include a small biological component<sup>34</sup>

<sup>k</sup> Dust aerosol optical depth (AOD) feedback strength estimate taken from Thornhill et al.<sup>36</sup> Naik et al.<sup>37</sup> also list a central estimate of -0.004 W/m<sup>2</sup>/K, but with “very likely range” (-0.02 to 0.01) W/m<sup>2</sup>/K.

<sup>l</sup> The feedback strength associated with sea-salt emissions alone was estimated by Paulot et al.<sup>39</sup> to be -0.08 W/m<sup>2</sup>/K and, more recently, by Naik et al.<sup>37</sup> to be -0.049 W/m<sup>2</sup>/K with a “very likely range” of (-0.13 to 0.03) W/m<sup>2</sup>/K.

<sup>m</sup> Ocean stratification feedbacks involve both abiotic and biological processes. Here, we consider only the abiotic aspects. The biological components are treated separately in the “ocean biological pump” feedback.

---

<sup>n</sup> Decreasing carbon uptake by the ocean occurs due to at least two abiotic processes. First, increasing ocean stratification means that water near the surface stays warm, reducing CO<sub>2</sub> absorption (see also “solubility pump” loop). Second, reduced ocean mixing results in less CO<sub>2</sub> from the atmosphere being buried under the ocean’s surface. See Mann<sup>41</sup> for a non-technical overview.

<sup>o</sup> We have not classified this feedback loop as being exceptionally “slow” (in its entirety), although some aspects – notably, potential AMOC collapse and recovery – operate on very long time scales. Note that large-scale overturning circulation together with ocean heat uptake can be viewed as a significant negative feedback.<sup>48,20</sup>

<sup>p</sup> Other climate feedbacks involving ocean circulation are also possible.<sup>20</sup> For example, the Walker circulation could intensify, leading to a negative feedback (due to the ocean thermostat mechanism) or weaken, leading to a positive feedback – see section 3.4.2 of Heinze et al. for details.<sup>20</sup>

<sup>q</sup> In terms of radiative change per degree of warming (W/m<sup>2</sup>/K), the effect of changing surface temperatures due to the slowdown of the AMOC falls under the Planck, water vapor, lapse rate, surface albedo (sea ice, etc.), and cloud feedbacks, depending on how it alters the pattern of warming. So the short-term changes in the AMOC are already included in these other feedbacks from this perspective.

<sup>r</sup> There may also be a feedback involving abiotic carbonate precipitation.<sup>51</sup>

<sup>s</sup> Ciais et al.<sup>50</sup> estimated this feedback strength in terms of CO<sub>2</sub> concentration at 4%/K. We converted this estimate to units of W/m<sup>2</sup>/K by multiplying it by the CO<sub>2</sub> forcing for a 1% rise, 0.04 W/m<sup>2</sup>/%, yielding 0.16 W/m<sup>2</sup>/K.

<sup>t</sup> Current evidence for methane derived from hydrates entering the atmosphere is limited and there may be mitigating factors in terms of a potential feedback loop<sup>55,58</sup>

<sup>u</sup> Range of estimates for net feedback strength presented in Table 1 of Klocke et al.<sup>59</sup>

<sup>v</sup> Although this feedback loop appears similar to the “glacier and ice sheet albedo” feedback loop, it differs in that the underlying mechanism is the lapse rate (elevation-temperature) relationship, which causes decreasing glacier and ice sheet elevations to result in increasing surface warming rates.<sup>23,60–62</sup>

<sup>w</sup> This is actually a set of three potential negative feedbacks that collectively act to increase the growth rate of thin ice – see Heinze et al.<sup>20</sup> for details. In addition, there are two other ice-ocean feedbacks in the Southern Ocean: the ice production–entrainment feedback (negative) and the ice-production-ocean–ocean-heat-storage feedback (positive).<sup>20,65</sup>

<sup>x</sup> The climate-ozone feedback involves many different processes that are difficult to cover in a simple table; for a detailed overview, see the “Climate-ozone feedback” subsection in section 6.4.5 of Naik et al.<sup>37</sup>

<sup>y</sup> Net ozone feedback and “very likely range of feedback parameter” estimated by Naik et al.<sup>37</sup> Note that changing lightning NO<sub>x</sub> emissions – sometimes treated as a separate feedback loop – can affect ozone production.<sup>36</sup> The lightning feedback loop strength was estimated by Naik et al.<sup>37</sup> to be -0.010 W/m<sup>2</sup>/K with a “very likely range” of -0.04 W/m<sup>2</sup>/K to 0.02 W/m<sup>2</sup>/K.

<sup>z</sup> We have flagged this loop with a dagger because it includes “climate CH<sub>4</sub> lifetime,” which is listed in Fig. 6.20 of Stocker et al.<sup>69</sup> (2013)

<sup>aa</sup> This feedback loop primarily involves water vapor, CH<sub>4</sub>, and O<sub>3</sub>.<sup>20</sup> Although we have classified it as “physical” (abiotic), it can be indirectly linked to biological processes (e.g., wetlands methane emissions). Methane lifetime may also be affected by changing lightning NO<sub>x</sub> emissions – sometimes treated as a separate feedback loop.<sup>36</sup>

<sup>bb</sup> The methane lifetime component of this feedback loop has an estimated strength of -0.010 W/m<sup>2</sup>/K with “very likely range” (-0.12 to 0.06) W/m<sup>2</sup>/K – see Table 6.8 of Naik et al.<sup>37</sup>

<sup>cc</sup> Climate change could also lead to an increase in methane emissions from wetlands due to other mechanisms<sup>76</sup>

<sup>dd</sup> Estimate from Table 15 of Thornhill et al.<sup>36</sup>

<sup>ee</sup> This feedback may be primarily associated with northern (temperate/boreal) lakes because they contain most of Earth’s ice-free freshwater.<sup>79</sup>

<sup>ff</sup> There is some uncertainty regarding the existence of a possible tipping point associated with Amazon (and other) forest dieback.<sup>1,85–91</sup> This uncertainty is partly due to the complexity of the system and how multiple anthropogenic pressures (including climate change and deforestation) are co-occurring. While many climate models suggest a tipping point may not exist,<sup>87</sup> even the latest generation of models (CMIP6) do not include all relevant processes that could create climate-vegetation feedbacks.<sup>92</sup> See section 5.4.9.1.1 of IPCC<sup>2</sup> for a recent overview of the literature. For reference, IPCC<sup>2</sup> estimated a maximum release of tropical land carbon of 200 GtC over the course of this century.

---

<sup>gg</sup> We use “dieback” to refer to a transition from forest to a new ecosystem state (e.g., grassland). In contrast, wildfires and insect outbreaks may have only temporary effects on forests, so these are treated as different feedback loops.

<sup>hh</sup> The effects of climate change on boreal forests and other woody vegetation are complex; for example, forested area in boreal regions could increase in response to moderate warming, but shrink in response to extreme warming<sup>99</sup>

<sup>ii</sup> But, see Phoenix and Bjerke<sup>101</sup> for a discussion of possible “Arctic browning.”

<sup>jj</sup> There is also be a negative feedback due to surface roughness changes affecting evapotranspiration (see also “evapotranspiration” loop) and heat fluxes.<sup>20,102</sup> However, the positive albedo feedback likely dominates.<sup>20</sup>

<sup>kk</sup> In some cases, this feedback loop may be viewed as a subset of the “plant stress” feedback loop.

<sup>ll</sup> Wildfires and smoldering combustion in peatlands are addressed in the “peatlands” feedback loop

<sup>mmm</sup> Over longer time scales, this effect may be partially offset by the production of pyrogenic carbon (charcoal), which is relatively stable.<sup>110,111</sup>

<sup>nn</sup> This is only the direct radiative perturbation<sup>34</sup>

<sup>oo</sup> Estimate with very likely range of feedback parameter based on Naik et al.<sup>37</sup>

<sup>pp</sup> Because of its importance, permafrost treated separately as the “permafrost” feedback loop

<sup>qq</sup> Numerous mechanisms are involved, including soil organic matter decomposition and microbial respiration<sup>113–117</sup>

<sup>rr</sup> This loop may have a human or social component since agricultural nitrogen fertilizer use is a primary source of N<sub>2</sub>O. Increasing N<sub>2</sub>O losses with a warming climate may leads to increasing demand for nitrogen fertilizer, which further increases N<sub>2</sub>O loss. Note that there are also several cooling effects related to nitrogen.<sup>121,122</sup>

<sup>ss</sup> Estimate from Xu-Ri et al.<sup>118</sup>; may be highly uncertain. More generally, Canadell et al.<sup>108</sup> estimated the total land N<sub>2</sub>O feedback strength is estimated to be  $0.02 \pm 0.01 \text{ W/m}^2/\text{K}$  and the total ocean N<sub>2</sub>O feedback strength is estimated to be  $-0.008 \pm 0.002 \text{ W/m}^2/\text{K}$ .

<sup>tt</sup> Multiple permafrost-related mechanisms are likely involved. For example, in addition to methane produced by microbial decay, permafrost thaw can lead to the release of natural gas (thermogenic methane).<sup>133</sup>

<sup>uu</sup> This is the 5<sup>th</sup>-95<sup>th</sup> percentile range from Burke et al.<sup>125</sup>; this estimate is emissions scenario dependent.<sup>125</sup> Note that this loop is contained within the “soil carbon (other)” loop, and thus the feedback strengths of these loops are not independent. According to Canadell et al.<sup>108</sup>, there is “low confidence” associated with the permafrost feedback with regard to timing, magnitude, and the relative roles of carbon dioxide and methane.

<sup>vv</sup> This occurs in response to rising atmospheric CO<sub>2</sub> concentration, but the effect is at least partly offset by CO<sub>2</sub> fertilization (see “CO<sub>2</sub> fertilization” feedback loop).<sup>20</sup>

<sup>ww</sup> This feedback loop covers microbes that are not included in the “ocean metabolic rates,” “freshwater ecosystems,” “soil carbon (other),” or “permafrost” feedback loops.

<sup>xx</sup> Unlike the “forest dieback” feedback loop, this includes terrestrial plants of all types. Unlike the “desertification” loop, plant stress need not involve a shift to arid/desert land cover.

<sup>yy</sup> One possible driver of desertification is a potential northward shift of the West African Monsoon (WAM) (which is also relevant to Sahara-Sahel greening<sup>139</sup>). Specifically, if the WAM moves northward, likely so will the subtropical high pressure that creates the Sahara, further shifting northward over the Mediterranean basin and amplifying desertification there. Similar effects could occur in places like northern New Mexico (if the Chihuahuan and Sonoran deserts shift northward).

<sup>zz</sup> A pulse of CO<sub>2</sub> emissions associated with plant community die-off is likely to occur along the desertification front.

<sup>aaa</sup> This feedback loop is related to the more general “terrestrial net primary productivity (NPP)” loop through potentially enhanced water-use efficiency of photosynthesis due to atmospheric CO<sub>2</sub> enrichment (except during extreme hotter drought episodes).

<sup>bbb</sup> However, if the West African monsoon moves northward, desertification in the Mediterranean basin could result – see “desertification” feedback loop footnote for details.

<sup>ccc</sup> The effects of rising CO<sub>2</sub> on productivity may be partly (or fully) countered by effects of increasing temperature, etc. (see “plant stress” feedback loop); thus, the overall effect of climate change on NPP may be uncertain.<sup>83</sup>

<sup>ddd</sup> However, climate change could reduce the rates at which plants can absorb carbon from the atmosphere due to triose phosphate utilization (TPU)-limited photosynthesis.<sup>143</sup> Similarly, climate change could lead to increasing



---

plant respiration rates<sup>144</sup> (although this effect might be relatively weak<sup>145</sup>) and to increasing CH<sub>4</sub> and N<sub>2</sub>O emissions by cryptogamic covers.<sup>146</sup> Other effects such as increasing heat wave occurrence are also possible.<sup>147</sup>

<sup>eee</sup> This estimate is from Norby et al.<sup>141</sup>

<sup>fff</sup> Arctic phytoplankton may also contribute substantially to Arctic warming through direct warming – another potential positive feedback loop<sup>154</sup>

<sup>ggg</sup> Estimate is for ocean bacterial respiration<sup>57</sup>

<sup>hhh</sup> The ocean biological pump involves many interrelated processes and factors that may be sensitive to climate change, including phytoplankton growth, deoxygenation, seawater viscosity, and calcification.<sup>156,158</sup> Climate change driven coral bleaching<sup>159</sup> may also affect the ocean biological pump. Ocean metabolic rates are handled separately in the “ocean metabolic rates” feedback loop, although they may be considered part of the ocean biological pump.

<sup>iii</sup> As noted by Pörtner et al.<sup>157</sup>, it is difficult to determine whether the ocean biological pump will act as a net positive or negative climate feedback. Moreover, there is “low confidence” with regard to how the changing biological pump will affect the magnitude and sign of the ocean carbon feedback.<sup>108</sup>

<sup>jjj</sup> Ocean acidification could also increase N<sub>2</sub>O production rates, representing another potential feedback loop.<sup>164</sup> However, this might be restricted to certain regions, and the opposite effect is also possible.<sup>165,164</sup>

<sup>kkk</sup> The sign of this feedback appears to be uncertain. For example, a negative sign is reported in Ciais et al.<sup>50</sup> based on HadGEM2-ES<sup>166</sup>, whereas the simulations of Wang et al.<sup>163</sup> suggest there is a small positive feedback associated with DMS globally. More recently, the DMS feedback strength was given as  $0.005 \pm 0.006$  W/m<sup>2</sup>/K by Thornhill et al.<sup>36</sup> So, we have opted to list the sign as uncertain, although we provide a positive central estimate for the strength based on Naik et al.<sup>37</sup>

<sup>lll</sup> Assessed central estimate and very likely range of the feedback parameter as given in Table 6.8 of Naik et al.<sup>37</sup>

**Table S2. Summary of 15 potential human climate feedback loops.** The first column indicates the primary system or process involved in the loop. “Effect of climate change” describes how climate change alters the system (an active process) and “Effect on climate change” describes how the response of the system contributes to climate change. Subsequent columns indicate overall/net feedback direction, strength, and type respectively. References for strength estimates are listed in the first column (“Feedback name”) or in the footnotes (when the feedback has multiple references). When a complete feedback loop has (to the best of our knowledge) not been presented before it, is shown in blue. Sub-category refers only to the dominant type or types associated with the feedback loop. When the overall feedback direction is unknown, this is indicated with a “?” symbol. Note that these human loops may include biological and physical aspects. Some feedback loops may be highly uncertain or speculative; especially, those for which feedback strength estimates (or confidence intervals, etc.) were unavailable. This is a list of example human feedback loops and is therefore not intended to be exhaustive.

<b>Feedback name</b>	<b>Effect of climate change</b>	<b>Effect on climate change</b>	<b>+/-</b>	<b>Approximate strength</b>	<b>Type</b>
1. Climate-related disasters	Increasing forest fire <sup>106</sup> , tropical storm <sup>167</sup> , flood <sup>168</sup> , and coastal erosion frequency or intensity in some regions	Increasing carbon costs associated with rebuilding <sup>mmmm</sup> , but may be offset by more energy-efficient replacements and negative impacts on economic growth	?		Natural disasters
2. Human migration	Increasing uninhabitable area due to extreme heat <sup>169</sup> , sea level rise <sup>170</sup> , permafrost thaw <sup>171</sup> , coastal flooding <sup>170</sup> , and potential soil degradation <sup>172</sup>	Increasing migration and construction carbon costs, movement to more carbon-intensive areas; conversely, potentially smaller carbon footprints (increasing poverty), energy efficient construction	?		Movement
3. Human mobility <sup>173</sup>	Potentially increasing movement (vehicle miles travelled) due to warming	Increasing CO <sub>2</sub> emissions	+		Movement
4. Transport routes <sup>174–176</sup>	Changing transport routes (e.g., due to decreasing Arctic sea ice)	Changing CO <sub>2</sub> emissions	?		Movement
5. Energy demand <sup>177 nnn</sup>	Increasing global mean temperature	Changing GHG emissions due to changing energy demand (e.g., increasing energy needed to cool buildings but decreasing energy needed for heating)	?	Heating and cooling CO <sub>2</sub> emissions projected to rise from 0.8 Gt C (2000) to 2.2 Gt C (2100)	Energy

Feedback name	Effect of climate change	Effect on climate change	+/-	Approximate strength	Type
<b>6.</b> Agriculture <sup>178</sup> -180 ppp	Changing crop yields overall, <sup>qqq</sup> changing agricultural suitability, water supply and demand	Changing GHG emissions due to agriculture, deforestation-related emissions due to migration of agricultural areas <sup>rrr</sup>	?		Food/water
<b>7.</b> Coral reefs <sup>181-183</sup>	Increasing coral die-off, loss of associated fisheries and other ecosystem services	Potentially increasing carbon costs due to alternative food production or migration <sup>sss</sup>	+	Roughly 850 million people likely derive benefits from coral reef ecosystem services <sup>ttt</sup>	Food/water
<b>8.</b> Freshwater	Declining freshwater availability in certain regions <sup>184,185</sup>	Increasing CO <sub>2</sub> emissions due to increasing migration or desalination <sup>186 uuu</sup>	+		Food/water
<b>9.</b> Mitigation <sup>188,189</sup>	Increasing sense of urgency as climate change intensifies and damages increase <sup>vvv</sup>	Increasing mitigation (climate action) efforts <sup>www</sup> (e.g., increasing renewables <sup>xxx</sup> , carbon prices, natural climate solutions such as reforestation or afforestation, bio-energy) <sup>yy</sup>	-	Collectively, mitigation may be the largest human feedback loop	Psychological
<b>10.</b> Policy paralysis <sup>193</sup>	Climate change becomes an increasingly large policy issue	Larger policy issues can be more or less difficult to address <sup>zzz</sup>	+		Psychological
<b>11.</b> Economic growth <sup>188,194</sup>	Decreasing macroeconomic production and human consumption <sup>aaaa</sup>	Decreasing emissions, but potentially decreasing investments in capital-intensive renewables and other mitigation strategies	?		Political/economic
<b>12.</b> Economic disruption <sup>188</sup>	Increasing frequency of economic disruptions (natural disasters, crop failures, etc.)	Decreasing investment in climate mitigation and associated technologies (due to focusing on immediate concerns)	+		Political/economic
<b>13.</b> Political disruption <sup>188</sup>	Increasing political upheaval (e.g., due to mass migrations)	Increasing nationalism leading to decreasing international cooperation on climate mitigation	+		Political/economic

Feedback name	Effect of climate change	Effect on climate change	+/-	Approximate strength	Type
14. Geopolitics <sup>193</sup> bbbb	Increasing large and unequal climate impacts on countries	Increasing difficulty of global cooperation (leading to decreasing mitigation)	+		Political/ economic
15. Human conflict <sup>195</sup>	Increasing conflict (declining resources)	Increasing GHG emissions associated with militaries	+		Political/ economic

<sup>mmm</sup> However, the carbon costs of rebuilding are likely small relative to those associated with new construction due to rising global demand.

<sup>nnn</sup> Although we refer primarily to energy demand for heating and cooling, this feedback loop is much broader. For example, energy is also required to produce green technologies such as batteries.

<sup>ooo</sup> With respect to heating and cooling, increasing need for air conditioning and decreasing need for heating roughly cancel each other out – feedback loop is slightly negative in the short term (demand for heating is the dominant factor) and slightly positive in the long term (demand for air conditioning is the dominant factor)<sup>177</sup>

<sup>ppp</sup> Other feedback loops involving land use change are also possible. For example, climate change could lead to additional forestry at high latitudes, with consequences for surface albedo, carbon sequestration, and so on.

<sup>qqq</sup> This could occur due to multiple factors, including crop failures as a result of extreme extents such as drought, extreme heat, and insect outbreaks.

<sup>rrr</sup> Shifting agricultural areas (especially due to soil degradation) can also lead to human migration – see “human migration” feedback loop

<sup>sss</sup> Coastal ecosystem damage (loss of CO<sub>2</sub> sequestration) as a result of losing coastal protection is also possible, but this is addressed under the “coastal productivity” biological feedback loop.

<sup>ttt</sup> See Burke et al.<sup>181</sup> for details

<sup>uuu</sup> In addition, large-scale desalination produces concentrated brine, which can have negative impacts on the environment and wildlife<sup>187</sup>

<sup>vvv</sup> Conversely, there may be an increasing sense of hopelessness as climate change intensifies (leading to decreasing mitigation efforts), but this effect may be relatively weak. The public’s response to climate change may be partly dependent on whether fear-inducing representations are employed.<sup>190</sup> Additionally, the public’s sense of urgency may be limited due to shifting baselines with respect to “normal” conditions.<sup>191</sup>

<sup>www</sup> Note that climate policy and air pollution policies are often connected and may involve complex, non-linear feedback loops<sup>189</sup>

<sup>xxx</sup> The physical impacts of climate change can also impact renewable energy potential, resulting in other potential feedback loops<sup>192</sup>

<sup>yyy</sup> These mitigation efforts can have many climate-related effects such as changing CO<sub>2</sub> and short-lived pollutant emissions and changing surface albedo.<sup>189</sup> Other mitigation efforts are also possible; for example, increasing funding to develop new technologies for decarbonization.

<sup>zzz</sup> But, this need not necessarily be the case – see “mitigation” feedback loop for details


<sup>aaaa</sup> Although not addressed separately, human population growth could be relevant to this and other human feedback loops.

<sup>bbbb</sup> This loop is closely linked to the “mitigation” loop in that climate mitigation efforts are often a geopolitical challenge

**Table S3. Credits for feedback loop pictures (Figures 1, S1).** Columns indicate the loop name and number (see Tables 1, S1), source/credit, license (Public Domain or Creative Commons), approximate location, and image URL (as of May 25, 2021).

Name	#	Credit	License	Location	URL
Sea ice albedo	3	Patrick Kelley	CC BY 2.0	The Arctic	<a href="https://en.wikipedia.org/wiki/Sea_ice#/media/File:Arctic_ice.jpg">https://en.wikipedia.org/wiki/Sea_ice#/media/File:Arctic_ice.jpg</a>
Ice sheets	4	Hannes Grobe	CC BY-SA 2.5	Greenland's east coast	<a href="https://en.wikipedia.org/wiki/Ice_sheet#/media/File:Greenland-ice_sheet_hq.jpg">https://en.wikipedia.org/wiki/Ice_sheet#/media/File:Greenland-ice_sheet_hq.jpg</a>
Snow cover	6	Natalia_K ollegova	Pixabay License	Unknown	<a href="https://pixabay.com/photos/mountains-tundra-summer-snow-5359560/">https://pixabay.com/photos/mountains-tundra-summer-snow-5359560/</a>
Clouds	7	Doggo19292	Public Domain	Monterrey, Mexico	<a href="https://en.wikipedia.org/wiki/Cloud#/media/File:Monterrey_Mexico_Clouds.jpg">https://en.wikipedia.org/wiki/Cloud#/media/File:Monterrey_Mexico_Clouds.jpg</a>
Dust	8	Advanstra	CC BY-SA 3.0	Surfers Paradise Beach	<a href="https://en.wikipedia.org/wiki/2009_Australian_dust_storm#/media/File:20090923_-_Dust_Storm_-_Surfers_Paradise_(looking_south).JPG">https://en.wikipedia.org/wiki/2009_Australian_dust_storm#/media/File:20090923_-_Dust_Storm_-_Surfers_Paradise_(looking_south).JPG</a>
CH <sub>4</sub> hydrates	13	USGS	Public Domain	Seafloor of the northern Gulf of Mexico	<a href="https://en.wikipedia.org/wiki/Methane_clathrate#/media/File:Gas_hydrate_under_carbonate_rock.jpg">https://en.wikipedia.org/wiki/Methane_clathrate#/media/File:Gas_hydrate_under_carbonate_rock.jpg</a>
Chem. weathering	20	Bernard Spragg. NZ	Public Domain	Waipu Caves, Northland, New Zealand	<a href="https://www.flickr.com/photos/volvob12b/20125209969">https://www.flickr.com/photos/volvob12b/20125209969</a>
Peatlands	21	NASA's Earth Observatory	CC BY 2.0	Indonesia	<a href="https://commons.wikimedia.org/wiki/File:Seeing_Through_the_Smoky_Pall-Observations_from_a_Grim_Indonesian_Fire_Season_%2823451153146%29.jpg">https://commons.wikimedia.org/wiki/File:Seeing_Through_the_Smoky_Pall-Observations_from_a_Grim_Indonesian_Fire_Season_%2823451153146%29.jpg</a>
Wetlands	22	Abrget47j	CC BY-SA 3.0	Viru Bog, Estonia	<a href="https://en.wikipedia.org/wiki/Viru_Bog#/media/File:Sunrise_at_viru_bog.jpg">https://en.wikipedia.org/wiki/Viru_Bog#/media/File:Sunrise_at_viru_bog.jpg</a>
Freshwater	23	Aaron Rees	CC BY-SA 4.0	West Coast region of New Zealand	<a href="https://en.wikipedia.org/wiki/Lake#/media/File:Lake_Kanier_e.jpg">https://en.wikipedia.org/wiki/Lake#/media/File:Lake_Kanier_e.jpg</a>
Forest dieback	24	NASA/JPL-Caltech		Western Brazil	<a href="https://www.nasa.gov/feature/jpl/nasa-finds-amazon-drought-leaves-long-legacy-of-damage">https://www.nasa.gov/feature/jpl/nasa-finds-amazon-drought-leaves-long-legacy-of-damage</a>
Northern greening	25	Logan Berner/ Northern Arizona University		Arctic tundra	<a href="https://climate.nasa.gov/news/3025/warming-temperatures-are-driving-arctic-greening/">https://climate.nasa.gov/news/3025/warming-temperatures-are-driving-arctic-greening/</a>
Insects	26	Jonhall	CC BY 3.0	E. C. Manning Provincial Park, British Columbia, Canada	<a href="https://en.wikipedia.org/wiki/Mountain_pine_beetle#/media/File:Pine_Beetle_in_Manning_Park.jpg">https://en.wikipedia.org/wiki/Mountain_pine_beetle#/media/File:Pine_Beetle_in_Manning_Park.jpg</a>
Wildfire	27	Nick-D	CC BY-SA 4.0	Viewed from Tuggeranong in southern Canberra, Australia	<a href="https://en.wikipedia.org/wiki/2019%E2%80%9320_Australian_bushfire_season#/media/File:Orroral_Valley_Fire_viewed_from_Tuggeranong_January_2020.jpg">https://en.wikipedia.org/wiki/2019%E2%80%9320_Australian_bushfire_season#/media/File:Orroral_Valley_Fire_viewed_from_Tuggeranong_January_2020.jpg</a>
Permafrost	31	Boris Radosavljevic	CC BY 2.0	Herschel Island, Canada	<a href="https://en.wikipedia.org/wiki/Permafrost#/media/File:Permafrost_in_Herschel_Island_001.jpg">https://en.wikipedia.org/wiki/Permafrost#/media/File:Permafrost_in_Herschel_Island_001.jpg</a>
Desertification	35	David Stanley	CC BY 2.0	Ennedi Mountains, Chad, Central Africa	<a href="https://en.wikipedia.org/wiki/Sahel#/media/File:Acacia_Trees_(24227057806).jpg">https://en.wikipedia.org/wiki/Sahel#/media/File:Acacia_Trees_(24227057806).jpg</a>
Coastal productivity	38	Kino Obusan	CC BY-SA 4.0	Puerto Princesa, Palawan, Philippines	<a href="https://en.wikipedia.org/wiki/Mangrove#/media/File:Mangroves_in_Puerto_Princesa_City.jpg">https://en.wikipedia.org/wiki/Mangrove#/media/File:Mangroves_in_Puerto_Princesa_City.jpg</a>
Ocean bio.	40	Alexander Vasenin	CC BY-SA 3.0	Manta Alley (near Komodo, Indonesia)	<a href="https://en.wikipedia.org/wiki/Bluestripe_snapper#/media/File:A_school_of_blue-striped_snappers_at_Manta_Alley.JPG">https://en.wikipedia.org/wiki/Bluestripe_snapper#/media/File:A_school_of_blue-striped_snappers_at_Manta_Alley.JPG</a>

**Supplemental Figures**




**Figure S1. Approximate feedback strengths.** Estimated strengths are shown for select physical and biological feedback loops. See Table S1 for details and uncertainty estimates (where available).



**Figure S2. Feedback loop pictures.** Loop descriptions and photo credits are as follows: Sea ice albedo - Sea ice melting or not forming → Decreasing albedo [Patrick Kelley, CC BY 2.0]; Ice sheets - Increasing glacier & ice sheet melting → Decreasing albedo [Hannes Grobe, CC BY-SA 2.5]; Snow cover - Snow metamorphosis and decreasing snow cover → Decreasing albedo [Natalia\_Kollegova, Pixabay License]; Clouds - Changing cloud distribution and optical properties → Changing cloud albedo and greenhouse effect [Doggo19292, Public Domain]; Dust - Changing dust aerosol abundance → Changing planetary albedo and greenhouse effect [Advanstra, CC BY-SA 3.0]; CH<sub>4</sub> hydrates - Increasing methane hydrate

dissociation rates → Increasing release of methane into the atmosphere [USGS, Public Domain]; Chem. Weathering - Increasing carbonate and silicate weathering rates → Increasing CO<sub>2</sub> taken out of the atmosphere [Bernard Spragg, NZ, Public Domain]; Peatlands - Decreasing soil organic carbon due to lowering of water table, increasing vulnerability to fire/smoldering → Increasing release of CO<sub>2</sub> into the atmosphere, decreasing carbon sequestration [NASA's Earth Observatory, CC BY 2.0]; Wetlands - Increasing precipitation and boreal near-surface soil moisture potentially leading to expansion of wetlands → Increasing CO<sub>2</sub> sequestration and methane emissions [Abrget47j, CC BY-SA 3.0]; Freshwater - Increasing aquatic plant growth rates and microbial methane production → Increasing methane emissions [Aaron Rees, CC BY-SA 4.0]; Forest dieback - Dieback of Amazon, boreal, and other forests → Loss of sequestration, change in albedo, decreasing evapotranspiration [NASA/JPL-Caltech]; Northern greening - Potential expansion of high latitude/elevation forests & woody vegetation into tundra → Increasing CO<sub>2</sub> sequestration, decreasing albedo [Logan Berner/ Northern Arizona University]; Insects - Changing insect distributions & abundances → Loss of sequestration, change in albedo [Jonhall, CC BY 3.0]; Wildfire - Increasing fire frequency and/or severity and/or extent → Increasing CO<sub>2</sub> emissions, loss of sequestration, change in albedo [Nick-D, CC BY-SA 4.0]; Permafrost - Increasing CO<sub>2</sub> and methane release → Increasing GHG emissions [Boris Radosavljevic, CC BY 2.0]; Desertification - Increasing chronic aridification & hotter drought stress extremes leading to expanding deserts → Decreasing CO<sub>2</sub> sequestration, increasing CO<sub>2</sub> emissions, and increasing albedo [David Stanley, CC BY 2.0]; Coastal productivity - Increasing degradation of coastal ecosystems and loss of coral reef protection → Decreasing carbon sequestration by coastal ecosystems [Kino Obusan, CC BY-SA 4.0]; Ocean bio. - Increasing CO<sub>2</sub> in ocean, ocean acidification, warming, decreasing upwelling → Changing effectiveness of ocean as a carbon sink, decreasing CO<sub>2</sub> uptake [Alexander Vasenin, CC BY-SA 3.0]. See Table S1 and S3 for more information on the feedback loops and photos respectively.





**Figure S3. Feedback loop diagrams.** The feedback loops operate as follows: (A) Arctic warming leads to melting sea ice, which leads to further warming because water has lower albedo (reflectance) than ice; (B) Increasing temperatures lead to permafrost thawing, which produces CO<sub>2</sub> and methane emissions, which in turn leads to further increasing temperatures; (C) Rising temperatures lead to increasing fire frequency or severity causing increasing CO<sub>2</sub> emissions, loss of sequestration, and changes in albedo, which causing more warming. See Table S1 for details. Photo credits: (A) Patrick Kelley, CC BY 2.0; (B) Boris Radosavljevic , CC BY 2.0; (C) Nick-D, CC BY-SA 4.0.

## Supplemental Experimental Procedures

### *Feedback loops*

Here, we describe the methods used to construct our comprehensive list of feedback loops.

We compiled an initial list of climate feedback loops by conducting a literature review using computerized searches of the Google Scholar and Web of Science databases. We considered both synthesis papers dealing with multiple feedback loops (e.g., Bony et al.<sup>8</sup>, Steffen et al.<sup>57</sup>, Sherwood et al.<sup>3</sup>) and papers concerned with single loops. We also selectively looked at references cited by these papers.

We grouped the feedback loops into three general categories: Physical (abiotic), Biological, and Human. Physical feedback loops (e.g., water vapor<sup>3</sup>) primarily involve abiotic systems, biological loops involve the biosphere in some way (e.g., forest dieback<sup>83</sup>), and human loops involve human or social systems (e.g., agriculture<sup>179</sup>). Note that biological loops may also involve physical components and human loops can involve (non-human) biological or physical components. We also identified feedback loop “types,” which correspond to subcategories. For example, physical feedback loops were divided into types including “surface albedo” and “ocean.” We used this “type” variable to group the feedback loops in our list (within the general categories).

Because of the complexity of human systems, the human feedback loops may be more speculative in nature and difficult to quantify. So, we present these potential feedback loops separately from the physical and biological loops. These human loops are examples only, and this list is not intended to be exhaustive. While not recognized in our categorization system, all loops are “Human” in the sense that they involve anthropogenic climate change.

The term “climate feedback” has been defined in a number of different ways (for details, see Bates et al.<sup>196</sup>). For our project, we were motivated by the definition of Stocker et al.<sup>197</sup>: “*An interaction mechanism between processes in the climate system is called a climate feedback when the result of an initial process triggers changes in a second process that in turn influence the initial one. A positive feedback intensifies the initial process and a negative feedback reduces it.*” However, we only considered feedbacks to global temperature, excluding internal feedbacks that may be covered under this definition (e.g., grass cover increasing fires, which increases grass cover). According to this definition, each feedback loop can be partitioned into two interacting processes. For example, warming in the Arctic causes ice to melt (the initial process) and melting ice in turn leads to further warming by decreasing albedo (the second process).<sup>8</sup> For each climate feedback loop, we identified these two processes, which we termed as the “effect of climate change” and the “effect on climate change.” We also categorized each feedback loop as positive, negative, or unknown/uncertain direction. Finally, we determined approximate estimates of feedback loop strengths where possible. Feedback strengths are quantified in a variety of ways, although units of  $W/m^2/K$  are commonly used. In some cases, only a rough proxy indicator of strength was available – for example, the number of people who likely benefit from coral-

related ecosystem services. When available, we included uncertainty estimates associated with feedback strengths (e.g., standard errors or confidence intervals).

While each feedback loop differs from the others in either the “effect of climate change,” or the “effect on climate change,” there may be overlaps among some of the feedback loops. Given that these feedback loops can involve many highly complex and interacting systems, we viewed some overlap as unavoidable. We used footnotes to indicate occurrences of partial overlap (e.g., between “forest dieback” and “plant stress” feedback loops). Our strength estimates are generally separate and additive, except in cases of overlapping feedback loops (e.g., permafrost and general soil carbon loop strengths are not additive because of overlap). In practice, combined strengths (e.g., for water vapor and lapse rate feedbacks together) may be preferable for modeling since they can be easier to estimate.<sup>59</sup>

In our feedback loops table, we included loops that vary in strength over time, many of which could eventually weaken. Likely examples include permafrost (finite capacity to emit greenhouse gases), sea ice (eventually there may be little or no sea ice), and forest dieback (eventually large forested areas could be converted to other ecosystem types, possibly halting this process).

The focus of our table is on feedback loops. However, climate tipping elements and tipping points are related concepts of major importance to the biosphere.<sup>1,198,199,46,200</sup> In particular, tipping elements (biophysical systems with multiple stable states that contribute to regulate the state of the planet) in the current Earth system are inherently (at least since we left the last Ice Age some 12,000 years ago) in a state dominated by negative feedbacks, i.e., which dampen warming. But, if tipping points are crossed, as far as we know today, some feedbacks will likely shift from negative to positive, acting to amplify warming. Consequently, we identified the feedback loops in our table that might involve tipping elements. We followed the framework of Lenton et al.<sup>1,46</sup>, and considered only the 15 “policy-relevant potential future tipping elements in the climate system” in Table 1 of Lenton et al.<sup>1</sup>

After constructing the preliminary tables of feedback loops, we had them reviewed by more than twenty climate feedback specialists (see main paper acknowledgments section). These specialists were typically authors of feedback loop papers that we cited. We invited them to propose modifications to our tables of feedback loops, including adding or removing loops, improving loop descriptions, or suggesting additional references to cite.

### *Remaining carbon budget*

We present information on the remaining carbon budget framework for context.<sup>201,202</sup> Specifically, we provide the estimate of Forster et al.<sup>203</sup> that the remaining carbon budget (relative to the start of 2023) associated with 1.5 °C warming is roughly 260 Gt CO<sub>2</sub>. This budget could be exhausted in just 6.5 years.<sup>203</sup> Note that this is lower than the associated estimate of 380 Gt CO<sub>2</sub> provided by the Global Carbon Project.<sup>204</sup>

## *Limitations*

We took an inclusive approach to identifying potential feedback loops, and thus some of the loops in our table may be speculative and potentially negligible. We have classified these loops as having uncertain or unknown direction in our table. For example, the dust feedback loop has an estimated strength ranging from -0.2 to 0.2 W/m<sup>2</sup> by 2100, making it potentially negligible.<sup>34</sup> In particular, feedback loops for which we could not obtain strength estimates or strength uncertainty estimates (e.g., confidence intervals) may be highly uncertain.

Some feedback loops (e.g., chemical weathering<sup>70</sup>) may have major impacts only on very long time scales. Where applicable, we have identified these loops in our table. Some tipping elements may be uncertain since tipping elements can be challenging to identify; see feedback loop footnotes for details.

Our table of feedback loops is intended to be a general, simplified overview of Earth's many climate feedback loops. Therefore, providing an accurate strength estimate, associated time scale (and time of occurrence), and level of uncertainty for each feedback loop is beyond the scope of our work. Most likely, some of the feedback loops in our table are already occurring whereas others are not yet occurring and may ultimately be relatively weak.

The references for each feedback loop that we provide are intended as a selection from the literature, rather than an exhaustive list. We have generally prioritized citing work that is relatively recent and comprehensive in nature. Thus, these references may be useful to readers looking for more information on specific feedback loops. In cases where some aspects of a feedback loop are controversial and actively being debated in the literature, our selection of references is not intended as an endorsement of any particular point of view.

## **Note S1. Feedback Loop Descriptions**

For each feedback loop, we provide a brief overview, typically using excerpts (at most 200 words) from the associated references (Tables S1, S2). Aside from citations, modifications to the quoted text (deletions or insertions) are indicated with square brackets. For consistency, citations within excerpts have been reformatted according to our numbering system. Feedback loops that, to the best of our knowledge, have never before been presented in their entirety are shown in blue. Loop names are followed by directions: positive (+), negative (-), uncertain (?). When available, pictures are provided below the loop names; the associated credits are given in Table S3. Explanatory notes are provided in purple text.

*Physical and biological feedback loops (Table S1)*

### **1. Planck (black body radiation) (-)**

“The Planck feedback represents the extra emission to space of [longwave] radiation arising from a vertically uniform warming of the surface and the atmosphere with no change in composition. Physical expectation for this feedback is that  $\lambda_{\text{Planck}} \approx -4\epsilon\sigma T^3 \approx -3.3 \text{ W m}^{-2} \text{ K}^{-1}$  for present-day conditions, and the values shown in [...] from [general circulation models] of  $-3.2 \pm 0.04 \text{ W m}^{-2} \text{ K}^{-1}$  (1-sigma)<sup>205–207</sup> and those from observations of interannual variability<sup>208</sup> are both in general agreement with this physical expectation.”<sup>3</sup>

### **2. Water vapor (+)**

“Water vapor absorption is strong across much of the longwave spectrum, generally with a logarithmic dependence on concentration. Additionally, the Clausius–Clapeyron equation describes a quasi-exponential increase in the water vapor–holding capacity of the atmosphere as temperature rises. Combined, these facts predict a strongly positive water vapor feedback providing that the water vapor concentration remains at roughly the same fraction of the saturation specific humidity (i.e., unchanged relative humidity). Indeed, the global warming associated with a carbon dioxide doubling is amplified by nearly a factor of 2 by the water vapor feedback considered in isolation from other feedbacks<sup>209</sup>, and possibly by as much as a factor of 3 or more when interactions with other feedbacks are considered.”<sup>210,8</sup>

“[A] warmer climate increases stratospheric water vapor, and because stratospheric water vapor is itself a greenhouse gas, this leads to further warming. An estimate of its magnitude from a climate model yields a value of  $+0.3 \text{ W}/(\text{m}^2 \cdot \text{K})$ , suggesting that this feedback plays an important role in our climate system.”<sup>7</sup>

### 3. Sea ice (+)



“There are a number of important feedbacks associated with sea ice that influence projected climate sensitivity. The net effect is such that changes in sea ice contribute to a projected amplification of climate warming in the Arctic region (e.g., Holland and Bitz<sup>211</sup>; Rind et al.<sup>212</sup>). They also contribute to the global mean warming. For example, Rind et al. (1995) showed that 20%–40% of the simulated global surface air temperature increase at 2xCO<sub>2</sub> conditions was associated with changes in the ice cover.”<sup>8</sup>

“Arguably the most important sea ice feedback is the influence of the ice area and surface state on the surface albedo. As sea ice melts under a climate-warming scenario, the highly reflective surface is lost, allowing increased solar absorption. This enhances the initial warming perturbation, resulting in a positive feedback.”<sup>8</sup>

“Sea ice also affects the surface energy budget by insulating the overlying atmosphere from the relatively warm ocean. As such, the extent and thickness of sea ice modifies the turbulent heat fluxes at the surface.”<sup>8</sup>

### 4. Greenland and ice sheet albedo (+)



“On longer timescales [...], a tipping point (when the ice sheet enters a state of irreversible mass loss and complete melting is initiated) exists as part of the coupled ice sheet–atmospheric system. This consists of two interrelated feedback mechanisms: the [surface mass balance]–elevation feedback, as

described above, and the melt–albedo feedback.<sup>213–215</sup> The latter acts on the surface energy balance, by allowing more absorption of solar radiation from a melting and darkening snow surface, or removal of all snow leading to a darker ice surface. This feedback may be enhanced by ice-based biological processes, such as the growth of algae.<sup>216</sup> Thus, the activation of these feedbacks can lead to self-sustained melting of the entire ice sheet, even if the anomalous climatic forcing is removed.”<sup>22</sup>

## 5. Sea level rise (+)

“Apart from the cryosphere, a small positive surface albedo feedback comes from the inundation of coastal lands by sea level rise which thus replaces land with a less reflective ocean surface. For the Last Glacial Maximum (LGM), the estimated radiative effect is of order  $1 \text{ W m}^{-2}$  (Köhler et al.<sup>217</sup>, see section 5.1). But because sea-level rise realized during 150 years and several K of warming would be limited to at most a few meters compared to the LGM change of over 100 meters, the resulting effective feedback is only of order  $0.01 \text{ W m}^{-2} \text{ K}^{-1}$ .”<sup>3</sup>

## 6. Snow cover (+)



“Snow-albedo feedback (SAF) enhances climatic anomalies in Northern Hemisphere [...] land masses because of two changes in the snowpack as surface air temperature [...] increases.<sup>8,28,218–228</sup> First, snow cover shrinks, and where it does it generally reveals a land surface that is much less reflective of solar radiation. Second, the remaining snow generally has a lower albedo due to snow metamorphosis. For example, wet melting snow, more common in a warmer climate, has a lower surface albedo than dry frozen snow<sup>229</sup>.”<sup>29</sup>

## 7. Clouds (+)



“The inferred positive total cloud feedback arises from several contributions. These include (1) a lifting of high cloud tops in warmer climates, as indicated by detailed numerical cloud simulations, observed trends since the 1980s, climate models and expected from theory; (2) a dissipation of tropical and midlatitude marine low cloud, probably due to increased mixing of environmental air into clouds as the climate warms, as indicated by observed cloud variability, and detailed numerical cloud simulations; and (3) a dissipation of warm-season low cloud over land due to decreasing boundary layer relative humidity, as expected because land warms faster than oceans, and as seen in observed humidity trends since the 1970s and in [general circulation model] simulations of warming. Meanwhile, a sizable negative feedback from clouds in tropical deep convection regions is inferred from observations of interannual variability but does not overwhelm the combined positive feedbacks from rising high cloud tops and reduced low-cloud cover. [...] Interannual fluctuations in [top-of-atmosphere] energy balance, which reflect the net effect of all cloud types, also point to a positive total feedback, suggesting that we have not missed any major feedbacks by assessing only a finite set of individual cloud types.”<sup>3</sup>

## 8. Dust (?)



“Atmospheric dust is also changing partly due to climate effects. For example, dust concentrations at Barbados show a four-fold increase since the 1960s driven by meteorological changes in the African source region.”<sup>230,34</sup>



“Changes in global dust deposition predicted by models are also highly variable, as are estimates of future changes in dust emissions, which may be positive or negative”<sup>34</sup>

“Wind speed, soil moisture and vegetation cover are climate-driven variables that strongly affect dust emission fluxes, size distribution and mineralogical composition and hence, indirectly, control dust transport, deposition and radiative effects. Transport is also controlled by the atmospheric circulation and wet deposition by precipitation. There is, consequently, clear potential for feedbacks involving dust associated with a changing climate. However, as in the case of emissions of biomass burning aerosols from wildfires, there are both natural and anthropogenic factors governing dust emissions.<sup>231</sup> The attribution of changing dust emissions to natural or human causes is very uncertain for this reason and also because, unlike for purely anthropogenic species, the choice of a base time to define natural emissions is essentially arbitrary.”<sup>34</sup>

## **9. Other aerosols (?)**

“A warmer climate could also affect the production and/or lifetime of aerosols, in particular, dust, sea salt, natural sources of SO<sub>2</sub>/SO<sub>4</sub> and reactive nitrogen species, and natural fires. Besides changes to the direct aerosol radiative effect,<sup>39</sup> this could lead to additional indirect aerosol effects on clouds<sup>32,232</sup> and fire-induced effects on surface albedo.”<sup>3</sup>

## **10. Ocean stratification (+)**

“In this study, we analysed changes in ocean stratification (using N<sup>2</sup>) in multiple datasets, finding consistent evidence for overall enhanced stratification in most regions of the world oceans, down to depths of 2,000 m.”<sup>40</sup>

“Increasing stratification has important climate implications. The expected decrease in ocean ventilation<sup>233,234</sup> could affect ocean heat and carbon uptake,<sup>235,236</sup> water mass formation<sup>233</sup> and tropical storm formation and strength.<sup>237</sup> The associated decrease in ocean mixing, moreover, is consistent with a decline in ocean oxygen concentration,<sup>234,238</sup> reduced nutrient flux<sup>239</sup> and alteration of marine productivity and biodiversity,<sup>236,239,240</sup> as observed.”<sup>40</sup>

“Warmer waters absorb less atmospheric carbon dioxide [...] Less ocean mixing also means that less of the atmospheric carbon dioxide gets buried beneath the ocean surface. So carbon pollution accumulates even faster in the atmosphere, causing yet more warming.”<sup>41</sup>

## **11. Ocean circulation (?)**

“While the Atlantic Meridional Overturning Circulation (AMOC) is projected to slow down under anthropogenic warming, the exact role of the AMOC in future climate change has not been fully quantified. [...] Our results show that a weakened AMOC can explain ocean cooling south of Greenland that resembles the North Atlantic warming hole and a reduced Arctic sea ice loss in all seasons with a


delay of about 6 years in the emergence of an ice-free Arctic in boreal summer. In the troposphere, a weakened AMOC causes an anomalous cooling band stretching from the lower levels in high latitudes to the upper levels in the tropics and [...].”<sup>47</sup>

“By [...], we can isolate the pattern of surface temperature change due to a weakened AMOC. We find that surface air temperature shows a ‘bipolar seesaw’ response,<sup>241–243</sup> with cooling in the Northern Hemisphere (NH) and warming in the Southern Hemisphere (SH)[...]. The largest cooling occurs south of Greenland in the North Atlantic and exceeds 3°C. This cooling seems related to a decreased northward heat transport induced by the weakened AMOC [...]. On a global scale, the weakened AMOC causes a 0.2°C cooling in global mean surface temperature by 2061–2080 [...].”<sup>47</sup>

## 12. Ocean solubility pump (+)

“The surface dissolution and equilibration of CO<sub>2</sub> with the atmosphere is well understood and quantified. It varies with the surface ocean conditions, in particular with temperature (solubility effect) and alkalinity. The capacity of the ocean to take up additional CO<sub>2</sub> for a given alkalinity decreases at higher temperature.”<sup>50</sup>

## 13. Ocean methane hydrates (+)



“Gas hydrate, a frozen, naturally-occurring, and highly-concentrated form of methane, sequesters significant carbon in the global system and is stable only over a range of low-temperature and moderate-pressure conditions. Gas hydrate is widespread in the sediments of marine continental margins and permafrost areas, locations where ocean and atmospheric warming may perturb the hydrate stability field and lead to release of the sequestered methane into the overlying sediments and soils. Methane and methane-derived carbon that escape from sediments and soils and reach the atmosphere could exacerbate greenhouse warming. The synergy between warming climate and gas hydrate dissociation feeds a popular perception that global warming could drive catastrophic methane releases [...] Methane hydrate is likely undergoing dissociation now on global upper continental slopes and on continental shelves that ring the Arctic Ocean. Many factors [...] mitigate the impact of gas hydrate dissociation on atmospheric greenhouse gas concentrations though. There is no conclusive

proof that hydrate-derived methane is reaching the atmosphere now, but more observational data and improved numerical models will better characterize the climate-hydrate synergy in the future.”<sup>55</sup>

#### **14. Lapse rate (+)**

“Variation with height of the temperature changes induced by an external climate forcing can also constitute a radiative feedback [...]. The tropospheric temperature lapse rate is controlled by radiative, convective, and dynamical processes. At extratropical latitudes, the lapse rate is constrained by baroclinic adjustment.<sup>244</sup> The temperature profile of deep convective atmospheres is nearly moist adiabatic,<sup>245</sup> and dynamical processes prevent the tropical atmosphere from maintaining substantial horizontal temperature gradients in the free troposphere. As a result, the temperature profile of the free troposphere is close to a moist adiabat throughout low latitudes.”<sup>8</sup>

“In response to global warming, at low latitudes [general circulation models] predict a larger tropospheric warming at altitudes than near the surface (consistent with the moist adiabatic stratification of the atmosphere), and thus a negative lapse rate feedback. At mid- and high latitudes, on the other hand, they predict a larger warming near the surface than at altitude (i.e., a positive lapse rate feedback). On average over the globe, the tropical lapse rate response dominates over the extratropical response, and the climate change lapse rate feedback is negative in most or all the [general circulation models] [...].”<sup>8</sup>

#### **15. Glacier and ice sheet elevation (+)**

“The next most well-characterized geometry/SMB [surface mass balance] positive feedback is the temperature-based elevation/SMB feedback<sup>246</sup> that depends on the presence of ablation areas and atmospheric temperature change with elevation (i.e., lapse rates). As a result of these factors, an initial externally forced increase in ablation—for example, due to increased summer temperature—will lower ablation area elevation. This in turn causes additional melting as the ice surface experiences warmer low-elevation temperatures. The same mechanism also operates in reverse (as with all feedback processes). The impact of this feedback increases with greater levels of elevation change: using idealized ice loss geometries in an atmospheric model, Hakuba et al.<sup>247</sup> estimated a  $\sim 2^{\circ}\text{C}$  increase in GrIS-averaged [Greenland ice sheet-averaged] surface temperatures for each 25% GrIS volume loss increment, which is similar to earlier studies (e.g., Ridley et al.<sup>248</sup>), with the majority of this temperature increase arising from temperature lapse rate considerations. The importance of the temperature-based height/SMB feedback may have played a critical role in glacial climates characterized by ice sheets with large ablation zones, by triggering rapid deglaciation as thresholds in ice sheet geometry initiated the height/SMB feedback across a massive ice sheet area.<sup>249,61</sup>

#### **16. Antarctic rainfall (+)**

“While we do not propose the [Miocene Climatic Optimum] Antarctica was ever completely ice-free, our results demonstrate any spatial retreat of the [Antarctic ice sheet] can increase precipitation, causing

associated warming of the deep ocean—changes perhaps having the ability both to accelerate ice melt of ice shelves and glaciers through hydrofracturing from increased precipitation falling into crevasses<sup>250,251</sup> and to accelerate ice melt of marine-based subglacial basins.<sup>251,252</sup> Although the temperature changes resulting from changing ice-sheet extent are similar to those resulting from CO<sub>2</sub> changes, our study does not include feedbacks to the carbon cycle or to the ice sheet itself, and therefore the significance of our results could be greater than indicated here. Our non-realistic sensitivity studies using only a skin thickness of ice demonstrate the importance of both surface albedo and roughness for a hydrologic control on [deep water temperature] evolution.”<sup>63</sup>

## 17. Sea ice growth rate (-)

“The fact that sea ice decrease is not self-accelerating<sup>253</sup> in the presence of the ice albedo feedback leads to the conclusion that negative sea ice feedbacks must exist. There are at least three potential mechanisms which lead to sea ice negative feedbacks. First, thinner ice is warmer and has a higher winter open-water fraction, which induces more [long-wave] emission. Second, thinner ice is less insulating. Third, thinner ice has less snow,<sup>254</sup> further decreasing the insulation power of the sea ice cover. Overall, these three mechanisms drastically (and non-linearly) increase the growth rate for thin ice<sup>64</sup> contributing to rapidly bringing sea ice back to its equilibrium thickness in response to a perturbation<sup>255</sup>. In the Southern Ocean, where the stratification of the water column is weaker than in the Arctic, two competing ice–ocean feedbacks have been documented.<sup>65</sup> The first feedback is negative and is termed ice production–entrainment feedback. [...] The second feedback is positive and termed ice-production–ocean-heat-storage feedback. [...]”<sup>20</sup>

## 18. Ozone (?)

“State-of-the-art climate models now include more climate processes which are simulated at higher spatial resolution than ever.<sup>256</sup> Nevertheless, some processes, such as atmospheric chemical feedbacks, are still computationally expensive and are often ignored in climate simulations.<sup>256,257</sup> [...] [We] find an increase in global mean surface warming of around 1°C (~20%) after 75 years when ozone is prescribed at pre-industrial levels compared with when it is allowed to evolve self-consistently in response to an abrupt 4×CO<sub>2</sub> forcing. [...] This has important implications for global model intercomparison studies<sup>256,257</sup> in which participating models often use simplified treatments of atmospheric composition changes that are neither consistent with the specified greenhouse gas forcing scenario nor with the associated atmospheric circulation feedbacks<sup>258–260</sup>.”<sup>66</sup>

“Tropospheric ozone shows a range of responses to climate with models generally agreeing that warmer climate will lead to decreases in the tropical lower troposphere owing to increased water vapour and increases in the sub-tropical to mid-latitude upper troposphere due to increases in lightning and stratosphere-to-troposphere transport.<sup>261</sup> A small positive feedback is estimated from climate-induced changes in global mean tropospheric ozone<sup>262</sup> while a small negative feedback is estimated by Heinze et al.<sup>20</sup> based on the model results of Stevenson et al.<sup>261,37</sup>

## 19. Chemical reaction rates (?)

“Changing reaction rates in atmospheric chemistry involving the dynamics of CH<sub>4</sub>, O<sub>3</sub>, and water vapour can lead to both positive and negative climate feedbacks [...]. The major removal process for CH<sub>4</sub> is reaction with the OH radical, but OH is in turn removed by CH<sub>4</sub>. This self-feedback has the effect of amplifying any changes to CH<sub>4</sub> production or removal by a factor  $f$ .<sup>263</sup> Parameter  $f$  depends on the change in the lifetime of CH<sub>4</sub> with changing CH<sub>4</sub> concentration.”<sup>20</sup>

“Increased water vapour leads to increased O<sub>3</sub> destruction. Model simulations show that globally the effect of enhanced O<sub>3</sub> loss due to increased water vapour dominates, so that O<sub>3</sub> concentrations in the free troposphere decrease in a warmer and wetter climate.<sup>264,265</sup> Stevenson et al.<sup>261</sup> found varying responses to climate change in different chemistry models. They all showed decreased O<sub>3</sub> over the oceans, but some showed increased O<sub>3</sub> over the tropical continents where NO<sub>x</sub> emissions from lightning increased, and some showed increased O<sub>3</sub> in the upper troposphere around the subtropical jets due to increased stratosphere–troposphere exchange.”<sup>20</sup>

## 20. Chemical weathering (-)



“On geologic time scales, the amount of CO<sub>2</sub> in the atmosphere is determined by processes such as organic-carbon and carbonate-carbon sedimentation and burial, carbonate, organic carbon, and silicate weathering on land, and volcanic and metamorphic release of CO<sub>2</sub> [...].<sup>266</sup> Many of these processes are sensitive to the state of the surface environment, including its temperature and the intensity of the hydrologic cycle. These environmental variables, in turn, are sensitive to atmospheric  $p\text{CO}_2$  through the greenhouse effect. Thus, it is reasonable to assume that there are negative feedback mechanisms at work over geologic time that stabilize atmospheric  $p\text{CO}_2$  and climate.”<sup>70</sup>

## 21. Peatlands (+)



“Globally, the amount of carbon stored in peats exceeds that stored in vegetation and is similar in size to the current atmospheric carbon pool. Fire is a threat to many peat-rich biomes and has the potential to disturb these carbon stocks. Peat fires are dominated by smouldering combustion [...]. In undisturbed peatlands, most of the peat carbon stock typically is protected from smouldering, and resistance to fire has led to a build-up of peat carbon storage in boreal and tropical regions over long timescales. But drying as a result of climate change and human activity lowers the water table in peatlands and increases the frequency and extent of peat fires. The combustion of deep peat affects older soil carbon that has not been part of the active carbon cycle for centuries to millennia, and thus will dictate the importance of peat fire emissions to the carbon cycle and feedbacks to the climate.”<sup>72</sup>

“Drying in peatlands also increases the depth of belowground fuel combustion, releasing carbon to the atmosphere that has been stored in soils for centuries to millennia, thus creating a positive feedback to the climate system.”<sup>72</sup>

## 22. Wetlands (precipitation) (+)



“The influence of temperature on CH<sub>4</sub> emissions, however, is also strongly dependent on hydrologic conditions.<sup>267</sup> Mean precipitation in already dry midlatitude and subtropical regions is predicted to decline under warming scenarios, whereas in the wet midlatitudes and northern wetland regions it is predicted to increase.<sup>268</sup> Extended droughts will lower water tables in wetlands and decrease CH<sub>4</sub>

emissions but increase CO<sub>2</sub> emissions.<sup>269</sup> Increased precipitation will raise water tables and even expand wetland areas thereby promoting C sequestration and also CH<sub>4</sub> emission. The balance between these two processes is critical in determining whether changes in wetlands are contributing to a positive or negative climate feedback. A rise in precipitation can also increase the rate of organic substrate leaching to deeper parts of the peat profile, leading to increased methanogenesis. CH<sub>4</sub> production rates at depth can be 2–4 times higher than in the top 1 m of peat, and this effect has been observed on decadal timescales.<sup>270</sup> Increased methanogenesis will put greater pressure on the oxidative capacity along the export pathway of this CH<sub>4</sub> whether vertically or laterally when dissolved in groundwater.”<sup>74</sup>

### 23. Freshwater ecosystems (+)



“Net emissions of the potent GHG methane from ecosystems represent the balance between microbial methane production (methanogenesis) and oxidation (methanotrophy), each with different sensitivities to temperature. How this balance will be altered by long-term global warming, especially in freshwaters that are major methane sources, remains unknown. Here we show that the experimental warming of artificial ponds over 11 years drives a disproportionate increase in methanogenesis over methanotrophy that increases the warming potential of the gases they emit. The increased methane emissions far exceed temperature-based predictions, driven by shifts in the methanogen community under warming, while the methanotroph community was conserved. Our experimentally induced increase in methane emissions from artificial ponds is, in part, reflected globally as a disproportionate increase in the capacity of naturally warmer ecosystems to emit more methane. Our findings indicate that as Earth warms, natural ecosystems will emit disproportionately more methane in a positive feedback warming loop.”<sup>80</sup>

## 24. Forest dieback (+)



Here, we focus on the effects of extreme heat and drought. Although wildfires and insect damage can also impact forests, we treated these as separate feedback loops because they often represent transient disturbances. We also distinguish “plant stress” (can include non-forest vegetation) from the overlapping “forest dieback” feedback loop.

“Greenhouse gas emissions have significantly altered global climate, and will continue to do so in the future. Increases in the frequency, duration, and/or severity of drought and heat stress associated with climate change could fundamentally alter the composition, structure, and biogeography of forests in many regions. Of particular concern are potential increases in tree mortality associated with climate-induced physiological stress and interactions with other climate-mediated processes such as insect outbreaks and wildfire. [...] Here we present the first global assessment of recent tree mortality attributed to drought and heat stress. Although episodic mortality occurs in the absence of climate change, studies compiled here suggest that at least some of the world's forested ecosystems already may be responding to climate change and raise concern that forests may become increasingly vulnerable to higher background tree mortality rates and die-off in response to future warming and drought, even in environments that are not normally considered water-limited. This further suggests risks to ecosystem services, including the loss of sequestered forest carbon and associated atmospheric feedbacks. [...] Overall, our review reveals the potential for amplified tree mortality due to drought and heat in forests worldwide.”<sup>82</sup>



## 25. Northern greening (+)



“Northern high latitude ecosystems are experiencing amplified climate warming<sup>271</sup> that will be exacerbated by a series of positive feedbacks,<sup>272</sup> the relative magnitudes of which remain uncertain. A terrestrial albedo feedback to climate is underway in these ecosystems and being driven by two processes: densification and northward expansion of woody vegetation<sup>273–275</sup> and changes in the extent and duration of snow cover<sup>272,276</sup>. Currently, snow melt advance exerts the strongest feedback on regional warming,<sup>272</sup> however, continued increases in tree and shrub cover will likely diminish its role<sup>272,277</sup>.”<sup>95</sup>

## 26. Insect outbreaks (forests) (+)



“The mountain pine beetle [...] is a native insect of the pine forests of western North America, and its populations periodically erupt into large-scale outbreaks.<sup>278–280</sup> During outbreaks, the resulting widespread tree mortality reduces forest carbon uptake and increases future emissions from the decay of killed trees. [...] Here we estimate that the cumulative impact of the beetle outbreak in the affected region during 2000–2020 will be 270 [Mt] carbon [...] Climate change has contributed to the unprecedented extent and severity of this outbreak<sup>6</sup>. Insect outbreaks such as this represent an important mechanism by which climate change may undermine the ability of northern forests to take up and store atmospheric carbon [...].”<sup>103</sup>

“Warming-induced [Siberian silkmoth (SSM)] outbreaks are one of the major driving factors of successions within the taiga zone. [...] We analyzed the migration of alpine and northerly SSM outbreak boundaries in Siberia and the impact of [...] climate variables and topography on the outbreak dynamics. [...] outbreak onset was promoted by increased dryness and active temperatures and decreased root zone moisture content in the spring-early summer period. [...] Climate warming contributes to SSM migration into former outbreak free conifer stands located in highlands and at northern latitudes.”<sup>104</sup>

## 27. Wildfire (+)



“Although certain biomes are sensitive to constraints on biomass productivity and others to atmospheric conditions promoting combustion, substantial and rapid shifts are projected for future fire activity across vast portions of the globe. In the near term, the most consistent increases in fire activity occur in biomes with already somewhat warm climates; decreases are less pronounced and concentrated primarily in a few tropical and subtropical biomes. However, models do not agree on the direction of near-term changes across more than 50% of terrestrial lands, highlighting major uncertainties in the next few decades. By the end of the century, the magnitude and the agreement in direction of change are projected to increase substantially. Most far-term model agreement on increasing fire probabilities (~62%) occurs at mid- to high-latitudes, while agreement on decreasing probabilities (~20%) is mainly in the tropics. Although our global models demonstrate that long-term environmental norms are very successful at capturing chronic fire probability patterns, future work is necessary to assess how much more explanatory power would be added through interannual variation in climate variables.”<sup>106</sup>

## 28. Biogenic volatile organic compounds (BVOCs) (+)

“[...] shows the possible climate feedbacks associated with biogenic [secondary organic aerosol (SOA)]. The main driver of the feedback is that climate exerts a strong control over the emission of BVOCs [...]. Increases in temperature are likely to lead to increased BVOC emissions and aerosol concentrations, resulting in increased aerosol radiative cooling and a potential negative feedback mechanism.”<sup>281,34</sup>

“Increased temperature causes increased BVOC emissions (e.g., Guenther et al.<sup>282</sup>) but may also reduce the partitioning of semi-volatile compounds to the particles. Increased temperature also modifies vegetation resulting in either further increased or decreased BVOC emissions.”<sup>34</sup>

“Increased CO<sub>2</sub> concentrations may directly inhibit leaf-level isoprene emission<sup>283,284</sup> but the fertilization effect of the CO<sub>2</sub> on plant growth can increase emission rates.”<sup>34</sup>

“The observational evidence for large-scale climate-driven changes in BVOC emissions and SOA formation is limited, unlike for wildfires [...]. In a good example of such a study, Palmer et al.<sup>285</sup> used a 6 year record of satellite-observed formaldehyde column to infer a 20–30% interannual variability in isoprene emission over the south-eastern United States driven primarily by variations in surface air temperature.”<sup>34</sup>

### **29. Soil carbon (other) (+)**

“The majority of the Earth’s terrestrial carbon is stored in the soil. If anthropogenic warming stimulates the loss of this carbon to the atmosphere, it could drive further planetary warming.<sup>286–289</sup> [...] By extrapolating [...] to the global scale, we provide estimates of soil carbon sensitivity to warming that may help to constrain Earth system model projections. Our empirical relationship suggests that global soil carbon stocks in the upper soil horizons will fall by  $30 \pm 30$  petagrams of carbon to  $203 \pm 161$  petagrams of carbon under one degree of warming, depending on the rate at which the effects of warming are realized. Under the conservative assumption that the response of soil carbon to warming occurs within a year, a business-as-usual climate scenario would drive the loss of  $55 \pm 50$  petagrams of carbon from the upper soil horizons by 2050. [...] Despite the considerable uncertainty in our estimates, the direction of the global soil carbon response is consistent across all scenarios. This provides strong empirical support for the idea that rising temperatures will stimulate the net loss of soil carbon to the atmosphere, driving a positive land carbon–climate feedback that could accelerate climate change.”<sup>116</sup>

### **30. Soil nitrous oxide (+)**

“Although fertilizer use is presumed to have greatly stimulated global N<sub>2</sub>O emission, nonagricultural (especially tropical) soils still represent a source of N<sub>2</sub>O equal to all of the anthropogenic sources combined.<sup>290</sup> This natural N<sub>2</sub>O source must be influenced by climate, as can be deduced from both field observations and manipulative experiments.<sup>291</sup> In particular, widespread enhancement of soil N<sub>2</sub>O emission by warming is to be expected, because both nitrification and denitrification are highly temperature-dependent processes with estimated optima as high as 38°C.<sup>292–295</sup> It has been proposed that the global soil source of N<sub>2</sub>O should increase as the atmosphere warms,<sup>296</sup> and that this climate-induced increase could even become as important for the global N<sub>2</sub>O budget as projected increases in the anthropogenic sources of N<sub>2</sub>O.”<sup>296,118</sup>

“The modelled temperature dependence of N<sub>2</sub>O emission (c.  $1 \text{ Tg N yr}^{-1} \text{ K}^{-1}$ ) implies a positive climate feedback which, over the lifetime of N<sub>2</sub>O (114 yr), could become as important as the climate–carbon cycle feedback caused by soil CO<sub>2</sub> release.”<sup>118</sup>

### 31. Permafrost (+)



Dissociation of methane hydrates, which can occur in permafrost, is treated separately under “methane hydrates.”

“Large quantities of organic carbon are stored in frozen soils (permafrost) within Arctic and sub-Arctic regions. A warming climate can induce environmental changes that accelerate the microbial breakdown of organic carbon and the release of the greenhouse gases carbon dioxide and methane. This feedback can accelerate climate change, but the magnitude and timing of greenhouse gas emission from these regions and their impact on climate change remain uncertain.”<sup>128</sup>

### 32. Evapotranspiration (+)

“Warming leads to an increase in evaporation from soils. This negative soil moisture anomaly leads to a positive surface temperature anomaly through the reduction in latent heat flux.<sup>297</sup> The result is a positive feedback. In addition to this physical feedback, there is a chemically forced feedback. Under rising atmospheric CO<sub>2</sub> concentrations, plants open their stomata (plant stomata; [...]) less widely [...].<sup>298,299</sup> This leads to a reduction in evapotranspiration over land, a decrease in latent heat flux, and respective warming. This overall positive feedback is somewhat reduced by a secondary negative feedback: CO<sub>2</sub> fertilization [...] will lead to an increase in carbon assimilation and a respective increase in LAI (leaf area index – area covered by leaf canopy in relation to ground area) and a slight increase in surface albedo.<sup>300</sup> An uncertainty associated with this feedback is the original underlying surface albedo (if this were high, then the feedback could even become reversed).”

### 33. Microbial respiration (other) (+)

“Understanding how the metabolic rates of prokaryotes respond to temperature is fundamental to our understanding of how ecosystem functioning will be altered by climate change, as these microorganisms are major contributors to global carbon efflux. Ecological metabolic theory suggests that species living at higher temperatures evolve higher growth rates than those in cooler niches due to thermodynamic constraints. Here, using a global prokaryotic dataset, we find that maximal growth rate

at thermal optimum increases with temperature for mesophiles (temperature optima  $\leq 45$  °C), but not thermophiles ( $\geq 45$  °C). Furthermore, short-term (within-day) thermal responses of prokaryotic metabolic rates are typically more sensitive to warming than those of eukaryotes. Because climatic warming will mostly impact ecosystems in the mesophilic temperature range, we conclude that as microbial communities adapt to higher temperatures, their metabolic rates and therefore, biomass-specific CO<sub>2</sub> production, will inevitably rise. Using a mathematical model, we illustrate the potential global impacts of these findings.”<sup>134</sup>

### **34. Plant stress (+)**

We treated “plant stress” as different from “forest dieback” because plant stress can apply to non-forest vegetation.

“While higher CO<sub>2</sub> concentrations may boost plant growth and simultaneously help plants conserve water, the co-occurrence of hot conditions during drought could exacerbate plant stress, and potentially lead to increased damage to tissues and higher rates of mortality.<sup>301–303</sup> This increase in plant stress could occur even if drought frequency remains constant and have major impacts on forest structure and functioning in a hotter world.”<sup>83,136</sup>

“Higher rates of mortality during drought could dramatically alter forest structure. The timescales associated with forest growth are significantly longer than those associated with death, such that increased mortality will reduce forest cover much faster than it can regenerate. [...] Loss of forest cover due to mortality increases the surface albedo and simultaneously decreases evapotranspiration rates and increases sensible heating.<sup>304</sup> [...] The relatively longer timescale of recovery from mortality could also lead to less plant cover, and thus less regulation by stomata of surface to atmosphere water flux. The mortality response to the hotter droughts expected in the future is a critical uncertainty in determining the impact of future droughts on forests, and is likely to be a major impact independent of any increase in drought frequency.”<sup>136</sup>

### 35. Desertification (+)



This feedback loop deals with climate change potentially leading to expanding deserts. However, the effects of climate change can also cause deserts to shrink in some cases (e.g., see “Sahara and Sahel greening”).

“[Desertification] typically results from the compound effect of climate change and land use. Changes in the global and regional patterns of precipitation can be major drivers of desertification and have historically led to the expansion and contraction of major deserts on Earth. In fact, while many of the existing deserts are very old and formed millions of years ago, most of them were affected by Pleistocene climatic changes and expanded at some point into areas that are currently much wetter (500–800 mm/yr). In those areas the temporary loss of vegetation cover can explain the formation of some of the sand seas that are currently stabilized by vegetation [74], including the Kalahari, southern Sahara, the High Plains (US), the Mega-Thar (India), the Kimberlies (Australia), and the Llanos and the Pampas (S. America). Thus, in the course of Earth’s history, several regions around the world experienced the alternation of wet and dry periods. It is interesting to analyze how climate has been changing in more recent times and whether climate change studies predict an expansion or contraction of arid lands on Earth.”<sup>137</sup>

### 36. Sahara and Sahel greening (+)

“In the future, the Sahara and Sahelian regions could experience more rainfall than today as a result of climate change. Wetter periods, termed African humid periods, occurred in the past and witnessed a mesic landscape in place of today’s hyperarid and semiarid environment. Such large past changes raise the question of whether the near future might hold in store similar environmental transformations, particularly in view of the growing human-induced climate, land-use, and land-cover changes.”<sup>139</sup>

### 37. CO<sub>2</sub> fertilization (+)

“One critical feedback occurs if C uptake by the biosphere increases in response to the fossil-fuel driven increase in atmospheric [CO<sub>2</sub>] (“CO<sub>2</sub> fertilization”), thereby slowing the rate of increase in atmospheric

[CO<sub>2</sub>]. Carbon exchanges between the terrestrial biosphere and atmosphere are often first represented in models as net primary productivity (NPP). However, the contribution of CO<sub>2</sub> fertilization to the future global C cycle has been uncertain, especially in forest ecosystems that dominate global NPP, and models that include a feedback between terrestrial biosphere metabolism and atmospheric [CO<sub>2</sub>] are poorly constrained by experimental evidence. We analyzed the response of NPP to elevated CO<sub>2</sub> (≈550 ppm) in four free-air CO<sub>2</sub> enrichment experiments in forest stands. We show that the response of forest NPP to elevated [CO<sub>2</sub>] is highly conserved across a broad range of productivity, with a stimulation at the median of 23 ± 2%. At low leaf area indices, a large portion of the response was attributable to increased light absorption, but as leaf area indices increased, the response to elevated [CO<sub>2</sub>] was wholly caused by increased light-use efficiency.”<sup>141</sup>

### 38. Coastal productivity (+)



“Mangroves will survive into the future but there have already been, and will continue to be, more negative than positive impacts due to climate change.”<sup>151</sup>

“As the world begins its transition to a low-carbon economy, removing atmospheric carbon dioxide (CO<sub>2</sub>) through biosequestration will be necessary to keep global warming under 2°C. Among the most efficient systems that provide biosequestration services are vegetated coastal habitats (VCHs), which include seagrasses, tidal marshes, and mangroves [...] and are known as ‘blue carbon’ ecosystems.<sup>149</sup> VCHs occupy only 0.2% of the ocean surface, yet contribute 50% of the total amount of carbon buried in marine sediments.<sup>150</sup> They have the ability to accumulate carbon without reaching saturation, and can store it in sediments over millennial timescales.<sup>149</sup> As with important terrestrial carbon sinks (eg Amazonian forests, permafrost regions), ecosystem degradation can shift blue carbon ecosystems from carbon sinks to carbon sources.<sup>305,152</sup>

### 39. Ocean metabolic rates (+)

The ocean represents a major sink for atmospheric carbon, partly through the ‘biological carbon pump’ in which sinking organic matter is exported from the surface to the deep sea.<sup>156</sup> The efficiency of the pump relies on a fine balance between the counteracting metabolic processes of carbon fixation into organic matter by photosynthesis and remineralization back to CO<sub>2</sub> by respiration.<sup>156</sup> When organic

carbon is moved to the deep ocean and/or buried, it creates a return flux from the atmosphere to the surface ocean. As the global ocean changes temperature, the balance between the two processes is likely to shift. If global temperatures increase, the efficiency of the ocean as a carbon sink should decline, and vice versa in a cooling world. This is potentially a very important positive feedback on global climate and atmospheric CO<sub>2</sub> levels.

#### 40. Ocean biological pump (?)



“Two ocean pumps play key roles in removing carbon from the surface ocean—the solubility pump is physico-chemically mediated whereas the biological pump is driven primarily by the interactions of marine biota from microbes to metazoa.<sup>306</sup> In brief, in the euphotic zone (Ez) photosynthetic carbon fixation by autotrophs, and heterotrophic bacterial production, inputs POC [particulate organic carbon] (and dissolved organic carbon [...]) to the biological pump. This POC supply to the upper ocean is subsequently modified and attenuated by a wide range of grazing activities which transform most of the phytoplankton and bacterial carbon into heterogeneous particles which eventually settle out of the surface ocean after a residence time of days to weeks.<sup>307,158</sup>”

“Pörtner et al.<sup>157</sup> in the Working Group 2 Fifth Assessment Report of the Intergovernmental Panel on Climate Change [...] concluded that due to the many “moving parts” it was not possible state with confidence how the sign or magnitude of the pump would alter in the coming decades. Instead they put forward a first assessment of the integrated knowledge platform (see Table 6.1, Pörtner et al.<sup>157</sup>) required to project how the biological pump will function in a future ocean.”<sup>158</sup>

#### 41. Phyto-plankton dimethyl sulfide (DMS) (+)

“Dimethyl sulfide (DMS) is a significant source of marine sulfate aerosol and plays an important role in modifying cloud properties. Fully coupled climate simulations using dynamic marine ecosystem and DMS calculations are conducted to estimate DMS fluxes under various climate scenarios and to examine the sign and strength of phytoplankton-DMS-climate feedbacks for the first time. Simulation results show small differences in the DMS production and emissions between pre-industrial and present climate scenarios, except for some areas in the Southern Ocean. There are clear changes in surface ocean DMS concentrations moving into the future, and they are attributable to changes in phytoplankton



production and competition driven by complex spatially varying mechanisms. Comparisons between parallel simulations with and without DMS fluxes into the atmosphere show significant differences in marine ecosystems and physical fields. Without DMS, the missing subsequent aerosol indirect effects on clouds and radiative forcing lead to fewer clouds, more solar radiation, and a much warmer climate.”<sup>163</sup>

## *Human feedback loops (Table S2)*

### **1. Climate-related disasters (?)**

In many regions of the world, climate change is projected to increase the frequency and severity of several types of climate-related disasters including forest fires<sup>106</sup>, tropical storms<sup>167</sup>, and floods<sup>168</sup>. These disasters can result in damages to buildings and infrastructure, which must be repaired or rebuilt. Since repairing and rebuilding have associated carbon costs, this constitutes a potential positive feedback loop wherein increasingly frequent or severe disasters results in increasing repairing and rebuilding carbon costs, which in turn contributes to further increasing disaster frequency or severity (through climate change). In 2009, 5.7 Gt of CO<sub>2</sub> emissions were associated with the global construction sector, representing 23% of all CO<sub>2</sub> emissions attributable to global economic activities<sup>308</sup>. Thus, rebuilding on a massive scale could have a substantial carbon footprint.

### **2. Human migration (?)**

In contrast to short-term disasters like hurricanes (see “climate-related disasters”) where in-place rebuilding may be an option, climate change can also render regions uninhabitable on a long-term basis. For example, sea level rise,<sup>170,309</sup> extreme mean annual temperatures,<sup>169</sup> and desertification<sup>137</sup> can all result in land becoming relatively unsuitable for human habitation. In such cases, humans may be forced to move and build new housing and infrastructure elsewhere. Given that both migration<sup>310,311</sup> and construction<sup>308</sup> can have significant carbon footprints, this could represent a significant positive feedback loop. Notably, sea level rise alone could force hundreds of millions of people to move by 2100.<sup>309</sup>

### **3. Human mobility (+)**

“Human behaviours alter—and are altered by—climate. Might the impacts of warming on human behaviours amplify anthropogenic contributions to climate change? Here, we show that warmer temperatures substantially increase transportation use in the USA. To do so, we combine meteorological data with data on vehicle miles travelled (VMT) and public transit trips (PTT) between 2002 and 2018. Moving from freezing temperatures up to 30°C increases VMT by over 10% and amplifies use of public transit by nearly 15%. Temperatures beyond 30°C exert little influence on either outcome. We then examine climate model projections to highlight the possible transportation impacts of future climatic changes. We project that warming over the coming century may add over one trillion cumulative VMT and six billion PTT in the USA alone, presenting the risk of a novel feedback loop in the human–environmental system.”<sup>173</sup>

### **4. Transport routes (?)**

“Rapid loss of sea ice is opening up the Arctic Ocean to shipping, a practice that is forecasted to increase rapidly by 2050 when many models predict that the Arctic Ocean will largely be free of ice toward the

end of summer. These forecasts carry considerable uncertainty because Arctic shipping was previously considered too sparse to allow for adequate validation. Here, we provide quantitative evidence that the extent of Arctic shipping in the period 2011–2014 is already significant and that it is concentrated (i) in the Norwegian and Barents Seas, and (ii) predominantly accessed via the Northeast and Northwest Passages. Thick ice along the forecasted direct trans-Arctic route was still present in 2014, preventing transit. Although Arctic shipping remains constrained by the extent of ice coverage, during every September, this coverage is at a minimum, allowing the highest levels of shipping activity. Access to Arctic resources, particularly fisheries, is the most important driver of Arctic shipping thus far.”<sup>176</sup>

## 5. Energy demand (?)

“In this article, we assess the potential development of energy use for future residential heating and air conditioning in the context of climate change. In a reference scenario, global energy demand for heating is projected to increase until 2030 and then stabilize. In contrast, energy demand for air conditioning is projected to increase rapidly over the whole 2000–2100 period, mostly driven by income growth. The associated CO<sub>2</sub> emissions for both heating and cooling increase from 0.8 Gt C in 2000 to 2.2 Gt C in 2100, i.e. about 12% of total CO<sub>2</sub> emissions from energy use (the strongest increase occurs in Asia). The net effect of climate change on global energy use and emissions is relatively small as decreases in heating are compensated for by increases in cooling. However, impacts on heating and cooling individually are considerable in this scenario, with heating energy demand decreased by 34% worldwide by 2100 as a result of climate change, and air-conditioning energy demand increased by 72%. At the regional scale considerable impacts can be seen, particularly in South Asia, where energy demand for residential air conditioning could increase by around 50% due to climate change, compared with the situation without climate change.”<sup>177</sup>

## 6. Agriculture (?)

There may also be climate feedbacks related to nitrous oxide emissions from fertilizer use.

“Change in yields, followed by a change in the area needed to supply the same amount of food. As discussed above, yields are projected to be affected from mildly positively to severely negatively, meaning that overall cropland will have to expand further into natural vegetation to compensate for somewhat lower average yields.”<sup>179</sup>

“Shifts in agricultural suitability, followed by relocation of agriculture. Cropping will follow agricultural suitability into higher latitudes, sometimes higher altitudes, causing deforestation of previously uncultivated areas and some reforestation on abandoned land.”<sup>179</sup>

“Changes in water demand for irrigated agriculture. In crop yield models, areas equipped with irrigation are ‘protected’ from any changes in precipitation and optimal evapotranspiration. That is, the models assume that enough water will be available for optimum irrigation. In reality, yields of irrigated agriculture can be affected if climate change alters the balance between the water available for

irrigation and irrigation water demand (it can influence both the supply and demand sides). This again would result in cropland expansion as compensation for lower yields.”<sup>179</sup>

## **7. Coral reefs (+)**

While coral reefs provide a number of important ecosystem services, especially with regard to food, income (e.g., from tourism), and storm protection, they are increasingly at risk due to climate change<sup>181-183</sup>. Thus, climate-related coral die-off could cause major problems for the approximately 850 million people who are likely to benefit from ecosystem services provided by coral reefs.<sup>181</sup> Many of these people may be forced to respond by migrating (with associated CO<sub>2</sub> emissions) and/or shifting to alternative means of food production that may have significant carbon costs. Because coral die-off could impact hundreds of million people, it is possible that this may be a significant positive feedback loop.

## **8. Freshwater (+)**

Climate change is projected to result in decreasing freshwater availability in some regions.<sup>184,185</sup> In these regions, humans may need to adapt by migrating (moving elsewhere),<sup>310,311</sup> scaling up desalination technology,<sup>186</sup> or importing freshwater. All of these options have associated carbon costs. Thus, declining freshwater availability could be part of a positive feedback loop where decreasing freshwater availability results in increasing greenhouse gas emissions.

## **9. Mitigation (-)**

Psychologically and behaviorally, there are two general ways in which humans might react to climate change becoming increasingly severe. Humans may respond by increasing mitigation efforts at the individual level (and higher levels). For example, by reducing air travel or shifting to plant-rich diets.<sup>312,313</sup> Conversely, some people may feel an increasing sense of hopelessness, apathy, or powerlessness,<sup>314</sup> leading them to decide that climate mitigation is futile and not worth pursuing. Depending on the relative strengths of these reactions, which could be affected by how climate change is represented,<sup>190</sup> “mitigation” could constitute a positive or negative feedback loop.

## **10. Policy paralysis (+)**

“Unlike most policy challenges, climate change gets worse the longer we take to address it.

How it works: The longer we wait to address climate change with major government action, the bigger the policy needed and the bigger economic impact that policy will have.

But the bigger the policy and economic hit get, the harder the politics get.

So we wait longer still, making the required policy and economic impact ever bigger, which makes the politics even more difficult.”<sup>193</sup>

## 11. Economic growth (?)

Note that climate-related effects on human population growth are also possible; this could form a related feedback loop.

“Empirical analysis of economic production trends generally finds a negative relationship between temperature and income.<sup>315</sup> Climate-related reduction in macroeconomic production will negatively affect human consumption, but may reduce emissions as well. The feedbacks associated with climate damages and growth are likely negative for the climate system, meaning that they lead to less severe climate change. This is because extreme climate outcomes become less probable as future emissions are dampened by the effects of current emissions on economic growth. But as noted by economist Martin Weitzman and others, this effect nevertheless would have profoundly harmful consequences for human well-being as future generations suffer substantial consumption losses compared to currently modeled damages.<sup>316,188</sup>

## 12. Economic disruption (+)

“The first pathway involves economic disruption caused by climate damages. Such disruptions are relatively easy to imagine: a sudden fall in agricultural productivity, the failure of critical infrastructure, or a string of high-impact natural disasters could all lead to severe economic disruptions that would result in a decline of national productivity. Given the interconnectedness of the global economic system, even if these harms did not befall the country in question, they could generate effects that propagated through the system, resulting in widespread costs. In the face of economic crisis, the attention of national leaders could turn from long-term global issues such as climate change to more pressing matters of economic stabilization. Investments in mitigation or adaptation might find themselves sacrificed for the needs of the day.”<sup>188</sup>

## 13. Political disruption (+)

“The second pathway involves political disruption caused by climate damages. For example, climate change-related events could lead to a wave of out migration from Bangladesh to nearby countries, causing political upheaval through an already unstable region. This climate change damage would turn into a positive self-reinforcing feedback if political leaders in India or China responded by embracing nationalist or isolationist positions or simply focused on the immediate crisis at hand, rather than long-term problems such as climate change. The basic relationships in these scenarios are between greenhouse gases, climate damages, economic or political disruption, and policy change.”<sup>188</sup>

## 14. Geopolitics (+)

“It takes global cooperation to address climate change, given its global nature. But climate change impacts different countries differently, so they're more likely to act on their own, and in their own self-interest.

But if there's no global cooperation, climate change continues to get worse — prolonging the adverse impacts on different countries, and giving them even less incentive to cooperate with other countries and more incentive to act on their own.”<sup>193</sup>

### **15. Human conflict (+)**

“According to Dr. Neta C. Crawford, Department Chair of Boston University’s Department of Political Science and the co-director of the study group Costs of War, military aggression and preparation exacerbates environmental problems that could lead to greater security risks and more war in the future as natural resources are depleted, causing a global refugee crisis.

‘The Pentagon is very worried about the stresses of climate change leading to displacement... and they’re concerned about climate war,’ Crawford tells Big Think in an interview. ‘They believe that it’s coming to a neighborhood near you.’

The problem, she notes, is that the Pentagon is a huge emitter of greenhouse gases and perpetrator of environmental destruction that increases the probability of war.”<sup>195</sup>

## Supplemental References

1. Lenton, T.M., Held, H., Kriegler, E., Hall, J.W., Lucht, W., Rahmstorf, S., and Schellnhuber, H.J. (2008). Tipping elements in the Earth's climate system. *Proceedings of the national Academy of Sciences* *105*, 1786–1793.
2. IPCC (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.
3. Sherwood, S., Webb, M.J., Annan, J.D., Armour, K., Forster, P.M., Hargreaves, J.C., Hegerl, G., Klein, S.A., Marvel, K.D., Rohling, E.J., et al. (2020). An assessment of Earth's climate sensitivity using multiple lines of evidence. *Reviews of Geophysics* *58*, e2019RG000678.
4. Forster, P., Storelvmo, T., Armour, K., Collins, W., J. L. Dufresne, D. Frame, D. J. Lunt, T. Mauritsen, M. D. Palmer, M. Watanabe, et al. (2021). The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment C Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. AZhai, A. Pirani, S. L. E Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.
5. Dessler, A., Zhang, Z., and Yang, P. (2008). Water-vapor climate feedback inferred from climate fluctuations, 2003–2008. *Geophysical Research Letters* *35*.
6. Romps, D.M., and Kuang, Z. (2009). Overshooting convection in tropical cyclones. *Geophysical Research Letters* *36*.
7. Dessler, A., Schoeberl, M., Wang, T., Davis, S., and Rosenlof, K. (2013). Stratospheric water vapor feedback. *Proceedings of the National Academy of Sciences* *110*, 18087–18091.
8. Bony, S., Colman, R., Kattsov, V.M., Allan, R.P., Bretherton, C.S., Dufresne, J.-L., Hall, A., Hallegatte, S., Holland, M.M., Ingram, W., et al. (2006). How well do we understand and evaluate climate change feedback processes? *Journal of Climate* *19*, 3445–3482.
9. Winton, M. (2006). Does the Arctic sea ice have a tipping point? *Geophysical Research Letters* *33*.
10. Hudson, S.R. (2011). Estimating the global radiative impact of the sea ice–albedo feedback in the Arctic. *Journal of Geophysical Research: Atmospheres* *116*.
11. Pistone, K., Eisenman, I., and Ramanathan, V. (2014). Observational determination of albedo decrease caused by vanishing Arctic sea ice. *Proceedings of the National Academy of Sciences* *111*, 3322–3326.
12. Cao, Y., Liang, S., Chen, X., and He, T. (2015). Assessment of sea ice albedo radiative forcing and feedback over the Northern Hemisphere from 1982 to 2009 using satellite and reanalysis data. *Journal of Climate* *28*, 1248–1259.

13. de Vernal, A., Hillaire-Marcel, C., Le Duc, C., Roberge, P., Brice, C., Matthiessen, J., Spielhagen, R.F., and Stein, R. (2020). Natural variability of the Arctic Ocean sea ice during the present interglacial. *Proceedings of the National Academy of Sciences* *117*, 26069–26075.
14. Thackeray, C.W., and Hall, A. (2019). An emergent constraint on future Arctic sea-ice albedo feedback. *Nature Climate Change* *9*, 972–978.
15. Chavas, J.-P., and Grainger, C. (2019). On the dynamic instability of Arctic sea ice. *npj Climate and Atmospheric Science* *2*, 1–7.
16. Tedesco, M., Mote, T., Fettweis, X., Hanna, E., Jeyaratnam, J., Booth, J., Datta, R., and Briggs, K. (2016). Arctic cut-off high drives the poleward shift of a new Greenland melting record. *Nature Communications* *7*, 1–6.
17. Hanna, E., Cropper, T.E., Hall, R.J., and Cappelen, J. (2016). Greenland Blocking Index 1851–2015: a regional climate change signal. *International Journal of Climatology* *36*, 4847–4861.
18. Liu, J., Chen, Z., Francis, J., Song, M., Mote, T., and Hu, Y. (2016). Has Arctic sea ice loss contributed to increased surface melting of the Greenland Ice Sheet? *Journal of Climate* *29*, 3373–3386.
19. Blackport, R., and Screen, J.A. (2020). Insignificant effect of Arctic amplification on the amplitude of midlatitude atmospheric waves. *Science advances* *6*, eaay2880.
20. Heinze, C., Eyring, V., Friedlingstein, P., Jones, C., Balkanski, Y., Collins, W., Fichet, T., Gao, S., Hall, A., Ivanova, D., et al. (2019). ESD Reviews: Climate feedbacks in the Earth system and prospects for their evaluation. *Earth System Dynamics* *10*, 379–452.
21. Naegeli, K., and Huss, M. (2017). Sensitivity of mountain glacier mass balance to changes in bare-ice albedo. *Annals of Glaciology* *58*, 119–129.
22. Pattyn, F., Ritz, C., Hanna, E., Asay-Davis, X., DeConto, R., Durand, G., Favier, L., Fettweis, X., Goelzer, H., Gollledge, N.R., et al. (2018). The Greenland and Antarctic ice sheets under 1.5 C global warming. *Nature Climate Change* *8*, 1053–1061.
23. Weertman, J. (1976). Milankovitch solar radiation variations and ice age ice sheet sizes. *Nature* *261*, 17–20.
24. Pattyn, F. (2018). The paradigm shift in Antarctic ice sheet modelling. *Nature communications* *9*, 1–3.
25. Sun, S., Pattyn, F., Simon, E.G., Albrecht, T., Cornford, S., Calov, R., Dumas, C., Gillet-Chaulet, F., Goelzer, H., Gollledge, N.R., et al. (2020). Antarctic ice sheet response to sudden and sustained ice-shelf collapse (ABUMIP). *Journal of Glaciology* *66*, 891–904.
26. Barron, E.J., Sloan II, J., and Harrison, C. (1980). Potential significance of land—sea distribution and surface albedo variations as a climatic forcing factor; 180 my to the present. *Palaeogeography, Palaeoclimatology, Palaeoecology* *30*, 17–40.



27. Nicholls, R.J., and Cazenave, A. (2010). Sea-level rise and its impact on coastal zones. *science* 328, 1517–1520.
28. Qu, X., and Hall, A. (2007). What controls the strength of snow-albedo feedback? *Journal of Climate* 20, 3971–3981.
29. Qu, X., and Hall, A. (2014). On the persistent spread in snow-albedo feedback. *Climate dynamics* 42, 69–81.
30. Thackeray, C.W., Qu, X., and Hall, A. (2018). Why do models produce spread in snow albedo feedback? *Geophysical Research Letters* 45, 6223–6231.
31. Trenberth, K.E., Zhang, Y., Fasullo, J.T., and Taguchi, S. (2015). Climate variability and relationships between top-of-atmosphere radiation and temperatures on Earth. *Journal of Geophysical Research: Atmospheres* 120, 3642–3659.
32. Gettelman, A., and Sherwood, S. (2016). Processes responsible for cloud feedback. *Current climate change reports* 2, 179–189.
33. Ceppi, P., Brient, F., Zelinka, M.D., and Hartmann, D.L. (2017). Cloud feedback mechanisms and their representation in global climate models. *Wiley Interdisciplinary Reviews: Climate Change* 8, e465.
34. Carslaw, K., Boucher, O., Spracklen, D., Mann, G., Rae, J., Woodward, S., and Kulmala, M. (2010). A review of natural aerosol interactions and feedbacks within the Earth system. *Atmospheric Chemistry & Physics* 10.
35. Kok, J.F., Ward, D.S., Mahowald, N.M., and Evan, A.T. (2018). Global and regional importance of the direct dust-climate feedback. *Nature Communications* 9, 1–11.
36. Thornhill, G., Collins, W., Olivié, D., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Gjermundsen, A., Horowitz, L., et al. (2021). Climate-driven chemistry and aerosol feedbacks in CMIP6 Earth system models. *Atmospheric Chemistry and Physics Discussions* 21, 1105–1126.
37. Naik, V., S. Szopa, B. Adhikary, P. Artaxo, T. Berntsen, W. D. Collins, S. Fuzzi, L. Gallardo, A. Kiendler Scharr, Z. Klimont, et al. (2021). Short-Lived Climate Forcers. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, J N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, B T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.
38. Thornhill, G.D., Collins, W.J., Kramer, R.J., Olivié, D., O’Connor, F., Abraham, N.L., Bauer, S.E., Deushi, M., Emmons, L., Forster, P., et al. (2020). Effective Radiative forcing from emissions of reactive gases and aerosols—a multimodel comparison. *Atmospheric Chemistry and Physics Discussions*, 1–29.
39. Paulot, F., Paynter, D., Winton, M., Ginoux, P., Zhao, M., and Horowitz, L.W. (2020). Revisiting the impact of sea salt on climate sensitivity. *Geophysical Research Letters* 47, e2019GL085601.

40. Li, G., Cheng, L., Zhu, J., Trenberth, K.E., Mann, M.E., and Abraham, J.P. (2020). Increasing ocean stratification over the past half-century. *Nature Climate Change*, 1–8.
41. Mann, M.E. (2020). The Oceans Appear to Be Stabilizing. Here’s Why it’s Very Bad News. *Newsweek*. <https://www.newsweek.com/climate-change-oceans-stabilizing-1534512>.
42. Buckley, M.W., and Marshall, J. (2016). Observations, inferences, and mechanisms of the Atlantic Meridional Overturning Circulation: A review. *Reviews of Geophysics* 54, 5–63.
43. Liu, W., Xie, S.-P., Liu, Z., and Zhu, J. (2017). Overlooked possibility of a collapsed Atlantic Meridional Overturning Circulation in warming climate. *Science Advances* 3, e1601666.
44. Williamson, M.S., Collins, M., Drijfhout, S.S., Kahana, R., Mecking, J.V., and Lenton, T.M. (2018). Effect of AMOC collapse on ENSO in a high resolution general circulation model. *Climate dynamics* 50, 2537–2552.
45. Gent, P.R. (2018). A commentary on the Atlantic meridional overturning circulation stability in climate models. *Ocean Modelling* 122, 57–66.
46. Lenton, T.M., Rockström, J., Gaffney, O., Rahmstorf, S., Richardson, K., Steffen, W., and Schellnhuber, H.J. (2019). Climate tipping points—too risky to bet against. *Nature* 575, 592–595.
47. Liu, W., Fedorov, A.V., Xie, S.-P., and Hu, S. (2020). Climate impacts of a weakened Atlantic Meridional Overturning Circulation in a warming climate. *Science advances* 6, eaaz4876.
48. Levitus, S., Antonov, J., Boyer, T., Garcia, H., and Locarnini, R. (2005). EOF analysis of upper ocean heat content, 1956–2003. *Geophysical research letters* 32.
49. Riebesell, U., Körtzinger, A., and Oschlies, A. (2009). Sensitivities of marine carbon fluxes to ocean change. *Proceedings of the National Academy of Sciences* 106, 20602–20609.
50. Ciais, P., Sabine, C., Bala, G., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., et al. (2014). Carbon and other biogeochemical cycles. In *Climate change 2013: the physical science basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge University Press), pp. 465–570.
51. Bialik, O.M., Sisma-Ventura, G., Vogt-Vincent, N., Silverman, J., and Katz, T. (2022). Role of oceanic abiotic carbonate precipitation in future atmospheric CO<sub>2</sub> regulation. *Scientific reports* 12, 1–8.
52. Shakhova, N., Semiletov, I., Salyuk, A., Yusupov, V., Kosmach, D., and Gustafsson, Ö. (2010). Extensive methane venting to the atmosphere from sediments of the East Siberian Arctic Shelf. *Science* 327, 1246–1250.
53. Skarke, A., Ruppel, C., Kodis, M., Brothers, D., and Lobecker, E. (2014). Widespread methane leakage from the sea floor on the northern US Atlantic margin. *Nature Geoscience* 7, 657–661.
54. Johnson, H.P., Miller, U.K., Salmi, M.S., and Solomon, E.A. (2015). Analysis of bubble plume distributions to evaluate methane hydrate decomposition on the continental slope. *Geochemistry, Geophysics, Geosystems* 16, 3825–3839.

55. Ruppel, C.D., and Kessler, J.D. (2017). The interaction of climate change and methane hydrates. *Reviews of Geophysics* 55, 126–168.
56. Shakhova, N., Semiletov, I., Gustafsson, O., Sergienko, V., Lobkovsky, L., Dudarev, O., Tumskey, V., Grigoriev, M., Mazurov, A., Salyuk, A., et al. (2017). Current rates and mechanisms of subsea permafrost degradation in the East Siberian Arctic Shelf. *Nature communications* 8, 1–13.
57. Steffen, W., Rockström, J., Richardson, K., Lenton, T.M., Folke, C., Liverman, D., Summerhayes, C.P., Barnosky, A.D., Cornell, S.E., Crucifix, M., et al. (2018). Trajectories of the Earth System in the Anthropocene. *Proceedings of the National Academy of Sciences* 115, 8252–8259.
58. Hong, W.-L., Torres, M.E., Carroll, J., Crémière, A., Panieri, G., Yao, H., and Serov, P. (2017). Seepage from an arctic shallow marine gas hydrate reservoir is insensitive to momentary ocean warming. *Nature communications* 8, 1–14.
59. Klocke, D., Quaas, J., and Stevens, B. (2013). Assessment of different metrics for physical climate feedbacks. *Climate dynamics* 41, 1173–1185.
60. Robinson, A., Calov, R., and Ganopolski, A. (2012). Multistability and critical thresholds of the Greenland ice sheet. *Nature Climate Change* 2, 429–432.
61. Fyke, J., Sergienko, O., Löfverström, M., Price, S., and Lenaerts, J.T. (2018). An overview of interactions and feedbacks between ice sheets and the Earth system. *Reviews of Geophysics* 56, 361–408.
62. Garbe, J., Albrecht, T., Levermann, A., Donges, J.F., and Winkelmann, R. (2020). The hysteresis of the Antarctic ice sheet. *Nature* 585, 538–544.
63. Bradshaw, C.D., Langebroek, P.M., Lear, C.H., Lunt, D.J., Coxall, H.K., Sostian, S.M., and de Boer, A.M. (2021). Hydrological impact of Middle Miocene Antarctic ice-free areas coupled to deep ocean temperatures. *Nature Geoscience*, 1–8.
64. Bitz, C., and Roe, G. (2004). A mechanism for the high rate of sea ice thinning in the Arctic Ocean. *Journal of Climate* 17, 3623–3632.
65. Goosse, H., Kay, J.E., Armour, K.C., Bodas-Salcedo, A., Chepfer, H., Docquier, D., Jonko, A., Kushner, P.J., Lecomte, O., Massonnet, F., et al. (2018). Quantifying climate feedbacks in polar regions. *Nature communications* 9, 1–13.
66. Nowack, P.J., Abraham, N.L., Maycock, A.C., Braesicke, P., Gregory, J.M., Joshi, M.M., Osprey, A., and Pyle, J.A. (2015). A large ozone-circulation feedback and its implications for global warming assessments. *Nature climate change* 5, 41–45.
67. Marsh, D.R., Lamarque, J.-F., Conley, A.J., and Polvani, L.M. (2016). Stratospheric ozone chemistry feedbacks are not critical for the determination of climate sensitivity in CESM1 (WACCM). *Geophysical Research Letters* 43, 3928–3934.
68. Holmes, C.D. (2018). Methane feedback on atmospheric chemistry: Methods, models, and mechanisms. *Journal of Advances in Modeling Earth Systems* 10, 1087–1099.

69. Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J., Nauels, A., Xia, Y., Bex, V., Midgley, P.M., et al. (2013). Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change 1535.
70. Kump, L.R., Brantley, S.L., and Arthur, M.A. (2000). Chemical weathering, atmospheric CO<sub>2</sub>, and climate. *Annual Review of Earth and Planetary Sciences* 28, 611–667.
71. Ise, T., Dunn, A.L., Wofsy, S.C., and Moorcroft, P.R. (2008). High sensitivity of peat decomposition to climate change through water-table feedback. *Nature Geoscience* 1, 763–766.
72. Turetsky, M.R., Benscoter, B., Page, S., Rein, G., Van Der Werf, G.R., and Watts, A. (2015). Global vulnerability of peatlands to fire and carbon loss. *Nature Geoscience* 8, 11–14.
73. Yang, G., Tian, J., Chen, H., Jiang, L., Zhan, W., Hu, J., Zhu, E., Peng, C., Zhu, Q., Zhu, D., et al. (2019). Peatland degradation reduces methanogens and methane emissions from surface to deep soils. *Ecological Indicators* 106, 105488.
74. Dean, J.F., Middelburg, J.J., Röckmann, T., Aerts, R., Blauw, L.G., Egger, M., Jetten, M.S., de Jong, A.E., Meisel, O.H., Rasigraf, O., et al. (2018). Methane feedbacks to the global climate system in a warmer world. *Reviews of Geophysics* 56, 207–250.
75. Zhang, Z., Zimmermann, N.E., Stenke, A., Li, X., Hodson, E.L., Zhu, G., Huang, C., and Poulter, B. (2017). Emerging role of wetland methane emissions in driving 21st century climate change. *Proceedings of the National Academy of Sciences* 114, 9647–9652.
76. Turetsky, M.R., Kotowska, A., Bubier, J., Dise, N.B., Crill, P., Hornibrook, E.R., Minkinen, K., Moore, T.R., Myers-Smith, I.H., Nykänen, H., et al. (2014). A synthesis of methane emissions from 71 northern, temperate, and subtropical wetlands. *Global change biology* 20, 2183–2197.
77. Pangala, S.R., Moore, S., Hornibrook, E.R., and Gauci, V. (2013). Trees are major conduits for methane egress from tropical forested wetlands. *New Phytologist* 197, 524–531.
78. Yvon-Durocher, G., Hulatt, C.J., Woodward, G., and Trimmer, M. (2017). Long-term warming amplifies shifts in the carbon cycle of experimental ponds. *Nature Climate Change* 7, 209–213.
79. Emilson, E.J., Carson, M.A., Yakimovich, K.M., Osterholz, H., Dittmar, T., Gunn, J., Myktyczuk, N., Basiliko, N., and Tanentzap, A. (2018). Climate-driven shifts in sediment chemistry enhance methane production in northern lakes. *Nature communications* 9, 1–6.
80. Zhu, Y., Purdy, K.J., Eyice, Ö., Shen, L., Harpenslager, S.F., Yvon-Durocher, G., Dumbrell, A.J., and Trimmer, M. (2020). Disproportionate increase in freshwater methane emissions induced by experimental warming. *Nature Climate Change* 10, 685–690.
81. Guo, M., Zhuang, Q., Tan, Z., Shurpali, N., Juutinen, S., Kortelainen, P., and Martikainen, P.J. (2020). Rising methane emissions from boreal lakes due to increasing ice-free days. *Environmental Research Letters* 15, 064008.

82. Allen, C.D., Macalady, A.K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D.D., Hogg, E.T., et al. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest ecology and management* 259, 660–684.
83. Allen, C.D., Breshears, D.D., and McDowell, N.G. (2015). On underestimation of global vulnerability to tree mortality and forest die-off from hotter drought in the Anthropocene. *Ecosphere* 6, 1–55.
84. Staal, A., Flores, B.M., Aguiar, A.P.D., Bosmans, J.H., Fetzer, I., and Tuinenburg, O.A. (2020). Feedback between drought and deforestation in the Amazon. *Environmental Research Letters* 15, 044024.
85. Nepstad, D.C., Stickler, C.M., Filho, B.S., and Merry, F. (2008). Interactions among Amazon land use, forests and climate: prospects for a near-term forest tipping point. *Philosophical transactions of the royal society B: biological sciences* 363, 1737–1746.
86. Nobre, C.A., and Borma, L.D.S. (2009). ‘Tipping points’ for the Amazon forest. *Current Opinion in Environmental Sustainability* 1, 28–36.
87. Huntingford, C., Zelazowski, P., Galbraith, D., Mercado, L.M., Sitch, S., Fisher, R., Lomas, M., Walker, A.P., Jones, C.D., Booth, B.B., et al. (2013). Simulated resilience of tropical rainforests to CO<sub>2</sub>-induced climate change. *Nature Geoscience* 6, 268–273.
88. Boulton, C.A., Good, P., and Lenton, T.M. (2013). Early warning signals of simulated Amazon rainforest dieback. *Theoretical Ecology* 6, 373–384.
89. Lovejoy, T.E., and Nobre, C. (2018). Amazon Tipping Point. *Science Advances* 4. 10.1126/sciadv.aat2340.
90. Amigo, I. (2020). When will the Amazon hit a tipping point? *Nature* 578, 505–508.
91. Walker, R.T. (2020). Collision course: Development pushes Amazonia toward its tipping point. *Environment: Science and Policy for Sustainable Development* 63, 15–25.
92. Sanderson, B.M., Pendergrass, A.G., Koven, C.D., Brient, F., Booth, B.B., Fisher, R.A., and Knutti, R. (2021). The potential for structural errors in emergent constraints. *Earth System Dynamics* 12, 899–918.
93. Jia, G.J., Epstein, H.E., and Walker, D.A. (2009). Vegetation greening in the Canadian Arctic related to decadal warming. *Journal of Environmental Monitoring* 11, 2231–2238.
94. Otto, J., Raddatz, T., and Claussen, M. (2011). Strength of forest-albedo feedback in mid-Holocene climate simulations. *Climate of the Past* 7, 1027–1039.
95. Lorant, M.M., Berner, L.T., Goetz, S.J., Jin, Y., and Randerson, J.T. (2014). Vegetation controls on northern high latitude snow-albedo feedback: observations and CMIP 5 model simulations. *Global change biology* 20, 594–606.

96. de Wit, H.A., Bryn, A., Hofgaard, A., Karstensen, J., Kvælev, M.M., and Peters, G.P. (2014). Climate warming feedback from mountain birch forest expansion: reduced albedo dominates carbon uptake. *Global Change Biology* 20, 2344–2355.
97. Thackeray, C.W., Fletcher, C.G., and Derksen, C. (2014). The influence of canopy snow parameterizations on snow albedo feedback in boreal forest regions. *Journal of Geophysical Research: Atmospheres* 119, 9810–9821.
98. Chae, Y., Kang, S.M., Jeong, S.-J., Kim, B., and Frierson, D.M. (2015). Arctic greening can cause earlier seasonality of Arctic amplification. *Geophysical Research Letters* 42, 536–541.
99. D’Orangeville, L., Houle, D., Duchesne, L., Phillips, R.P., Bergeron, Y., and Kneeshaw, D. (2018). Beneficial effects of climate warming on boreal tree growth may be transitory. *Nature communications* 9, 1–10.
100. Myers-Smith, I.H., Kerby, J.T., Phoenix, G.K., Bjerke, J.W., Epstein, H.E., Assmann, J.J., John, C., Andreu-Hayles, L., Angers-Blondin, S., Beck, P.S., et al. (2020). Complexity revealed in the greening of the Arctic. *Nature Climate Change* 10, 106–117.
101. Phoenix, G.K., and Bjerke, J.W. (2016). Arctic browning: extreme events and trends reversing arctic greening. *Global change biology* 22, 2960–2962.
102. Gustafsson, D., Lewan, E., and Jansson, P.-E. (2004). Modeling water and heat balance of the boreal landscape—comparison of forest and arable land in Scandinavia. *Journal of applied meteorology* 43, 1750–1767.
103. Kurz, W.A., Dymond, C., Stinson, G., Rampley, G., Neilson, E., Carroll, A., Ebata, T., and Safranyik, L. (2008). Mountain pine beetle and forest carbon feedback to climate change. *Nature* 452, 987–990.
104. Kharuk, V.I., Im, S.T., and Soldatov, V.V. (2020). Siberian silkmoth outbreaks surpassed geoclimatic barrier in Siberian Mountains. *Journal of Mountain Science* 17, 1891–1900.
105. Cudmore, T.J., Björklund, N., Carroll, A.L., and Staffan Lindgren, B. (2010). Climate change and range expansion of an aggressive bark beetle: evidence of higher beetle reproduction in naïve host tree populations. *Journal of Applied Ecology* 47, 1036–1043.
106. Moritz, M.A., Parisien, M.-A., Batllori, E., Krawchuk, M.A., Van Dorn, J., Ganz, D.J., and Hayhoe, K. (2012). Climate change and disruptions to global fire activity. *Ecosphere* 3, 1–22.
107. Liu, Y., Goodrick, S., and Heilman, W. (2014). Wildland fire emissions, carbon, and climate: Wildfire–climate interactions. *Forest Ecology and Management* 317, 80–96.
108. Canadell, J.G., G., J., Monteiro, P.M.S., Costa, M.H., Cunha, L.C. da, Cox, P.M., Eliseev, A.V., Henson, S., Ishii, M., Jaccard, S., et al. (2021). Global Carbon and other Biogeochemical Cycles and Feedbacks. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T. K. Maycock, T. Waterfield, O. Yelekçi, R. Yu and B. Zhou (eds.)]. Cambridge University Press. In Press.

109. Descals, A., Gaveau, D.L.A., Verger, A., Sheil, D., Naito, D., and Peñuelas, J. (2022). Unprecedented fire activity above the Arctic Circle linked to rising temperatures. *Science* 378, 532–537. 10.1126/science.abn9768.
110. Seiler, W., and Crutzen, P.J. (1980). Estimates of gross and net fluxes of carbon between the biosphere and the atmosphere from biomass burning. *Climatic change* 2, 207–247.
111. Lasslop, G., Coppola, A.I., Voulgarakis, A., Yue, C., and Veraverbeke, S. (2019). Influence of fire on the carbon cycle and climate. *Current Climate Change Reports* 5, 112–123.
112. Sporre, M.K., Blichner, S.M., Karset, I.H., Makkonen, R., and Berntsen, T.K. (2019). BVOC–aerosol–climate feedbacks investigated using NorESM. *Atmospheric Chemistry and Physics* 19, 4763–4782.
113. Ni, X., and Groffman, P.M. (2018). Declines in methane uptake in forest soils. *Proceedings of the National Academy of Sciences* 115, 8587–8590.
114. Conant, R.T., Ryan, M.G., \AAgren, G.I., Birge, H.E., Davidson, E.A., Eliasson, P.E., Evans, S.E., Frey, S.D., Giardina, C.P., Hopkins, F.M., et al. (2011). Temperature and soil organic matter decomposition rates—synthesis of current knowledge and a way forward. *Global Change Biology* 17, 3392–3404.
115. Karhu, K., Auffret, M.D., Dungait, J.A., Hopkins, D.W., Prosser, J.I., Singh, B.K., Subke, J.-A., Wookey, P.A., \AAgren, G.I., Sebastia, M.-T., et al. (2014). Temperature sensitivity of soil respiration rates enhanced by microbial community response. *Nature* 513, 81–84.
116. Crowther, T.W., Todd-Brown, K.E., Rowe, C.W., Wieder, W.R., Carey, J.C., Machmuller, M.B., Snoek, B., Fang, S., Zhou, G., Allison, S.D., et al. (2016). Quantifying global soil carbon losses in response to warming. *Nature* 540, 104–108.
117. Nottingham, A.T., Meir, P., Velasquez, E., and Turner, B.L. (2020). Soil carbon loss by experimental warming in a tropical forest. *Nature* 584, 234–237.
118. Xu-Ri, Prentice, I.C., Spahni, R., and Niu, H.S. (2012). Modelling terrestrial nitrous oxide emissions and implications for climate feedback. *New Phytologist* 196, 472–488.
119. Butterbach-Bahl, K., Baggs, E.M., Dannenmann, M., Kiese, R., and Zechmeister-Boltenstern, S. (2013). Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B: Biological Sciences* 368, 20130122.
120. Stocker, B.D., Roth, R., Joos, F., Spahni, R., Steinacher, M., Zaehle, S., Bouwman, L., Prentice, I.C., and others (2013). Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios. *Nature Climate Change* 3, 666–672.
121. Erisman, J.W., Galloway, J., Seitzinger, S., Bleeker, A., and Butterbach-Bahl, K. (2011). Reactive nitrogen in the environment and its effect on climate change. *Current Opinion in Environmental Sustainability* 3, 281–290.

122. Pinder, R.W., Davidson, E.A., Goodale, C.L., Greaver, T.L., Herrick, J.D., and Liu, L. (2012). Climate change impacts of US reactive nitrogen. *Proceedings of the National Academy of Sciences* *109*, 7671–7675.
123. Isaksen, I.S., Gauss, M., Myhre, G., Walter Anthony, K.M., and Ruppel, C. (2011). Strong atmospheric chemistry feedback to climate warming from Arctic methane emissions. *Global Biogeochemical Cycles* *25*.
124. Schuur, E.A., Abbott, B., Bowden, W., Brovkin, V., Camill, P., Canadell, J., Chanton, J., Chapin, F., Christensen, T., Ciais, P., et al. (2013). Expert assessment of vulnerability of permafrost carbon to climate change. *Climatic Change* *119*, 359–374.
125. Burke, E.J., Jones, C.D., and Koven, C.D. (2013). Estimating the permafrost-carbon climate response in the CMIP5 climate models using a simplified approach. *Journal of Climate* *26*, 4897–4909.
126. Schaefer, K., Lantuit, H., Romanovsky, V.E., Schuur, E.A., and Witt, R. (2014). The impact of the permafrost carbon feedback on global climate. *Environmental Research Letters* *9*, 085003.
127. Hodgkins, S.B., Tfaily, M.M., McCalley, C.K., Logan, T.A., Crill, P.M., Saleska, S.R., Rich, V.I., and Chanton, J.P. (2014). Changes in peat chemistry associated with permafrost thaw increase greenhouse gas production. *Proceedings of the National Academy of Sciences* *111*, 5819–5824.
128. Schuur, E.A., McGuire, A.D., Schädel, C., Grosse, G., Harden, J., Hayes, D.J., Hugelius, G., Koven, C.D., Kuhry, P., Lawrence, D.M., et al. (2015). Climate change and the permafrost carbon feedback. *Nature* *520*, 171–179.
129. Schädel, C., Bader, M.K.-F., Schuur, E.A., Biasi, C., Bracho, R., Čapek, P., De Baets, S., Diáková, K., Ernakovich, J., Estop-Aragones, C., et al. (2016). Potential carbon emissions dominated by carbon dioxide from thawed permafrost soils. *Nature climate change* *6*, 950–953.
130. Anthony, K.W., von Deimling, T.S., Nitze, I., Frohling, S., Emond, A., Daanen, R., Anthony, P., Lindgren, P., Jones, B., and Grosse, G. (2018). 21st-century modeled permafrost carbon emissions accelerated by abrupt thaw beneath lakes. *Nature communications* *9*, 1–11.
131. Knoblauch, C., Beer, C., Liebner, S., Grigoriev, M.N., and Pfeiffer, E.-M. (2018). Methane production as key to the greenhouse gas budget of thawing permafrost. *Nature Climate Change* *8*, 309–312.
132. Turetsky, M.R., Abbott, B.W., Jones, M.C., Anthony, K.W., Olefeldt, D., Schuur, E.A., Grosse, G., Kuhry, P., Hugelius, G., Koven, C., et al. (2020). Carbon release through abrupt permafrost thaw. *Nature Geoscience* *13*, 138–143.
133. Froitzheim, N., Majka, J., and Zastrozhnov, D. (2021). Methane release from carbonate rock formations in the Siberian permafrost area during and after the 2020 heat wave. *Proceedings of the National Academy of Sciences* *118*. 10.1073/pnas.2107632118.
134. Smith, T.P., Thomas, T.J., García-Carreras, B., Sal, S., Yvon-Durocher, G., Bell, T., and Pawar, S. (2019). Community-level respiration of prokaryotic microbes may rise with global warming. *Nature communications* *10*, 1–11.



135. Cavicchioli, R., Ripple, W.J., Timmis, K.N., Azam, F., Bakken, L.R., Baylis, M., Behrenfeld, M.J., Boetius, A., Boyd, P.W., Classen, A.T., et al. (2019). Scientists' warning to humanity: microorganisms and climate change. *Nature Reviews Microbiology* 17, 569–586.
136. Swann, A.L. (2018). Plants and drought in a changing climate. *Current Climate Change Reports* 4, 192–201.
137. D'Odorico, P., Bhattachan, A., Davis, K.F., Ravi, S., and Runyan, C.W. (2013). Global desertification: drivers and feedbacks. *Advances in water resources* 51, 326–344.
138. Shukla, P., Skea, J., Calvo Buendia, E., Masson-Delmotte, V., Pörtner, H., Roberts, D., Zhai, P., Slade, R., Connors, S., Van Diemen, R., et al. (2019). IPCC, 2019: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems.
139. Pausata, F.S.R., Gaetani, M., Messori, G., Berg, A., de Souza, D.M., Sage, R.F., and deMenocal, P.B. (2020). The Greening of the Sahara: Past Changes and Future Implications. *One Earth* 2, 235–250.
140. Bathiany, S., Claussen, M., and Brovkin, V. (2014). CO<sub>2</sub>-induced Sahel greening in three CMIP5 Earth system models. *Journal of Climate* 27, 7163–7184.
141. Norby, R.J., DeLucia, E.H., Gielen, B., Calfapietra, C., Giardina, C.P., King, J.S., Ledford, J., McCarthy, H.R., Moore, D.J., Ceulemans, R., et al. (2005). Forest response to elevated CO<sub>2</sub> is conserved across a broad range of productivity. *Proceedings of the National Academy of Sciences* 102, 18052–18056.
142. De Kauwe, M.G., Keenan, T.F., Medlyn, B.E., Prentice, I.C., and Terrer, C. (2016). Satellite based estimates underestimate the effect of CO<sub>2</sub> fertilization on net primary productivity. *Nature Climate Change* 6, 892–893.
143. Lombardozzi, D.L., Smith, N.G., Cheng, S.J., Dukes, J.S., Sharkey, T.D., Rogers, A., Fisher, R., and Bonan, G.B. (2018). Triose phosphate limitation in photosynthesis models reduces leaf photosynthesis and global terrestrial carbon storage. *Environmental Research Letters* 13, 074025.
144. Huntingford, C., Atkin, O.K., Martinez-De La Torre, A., Mercado, L.M., Heskell, M.A., Harper, A.B., Bloomfield, K.J., O'Sullivan, O.S., Reich, P.B., Wythers, K.R., et al. (2017). Implications of improved representations of plant respiration in a changing climate. *Nature Communications* 8, 1–11.
145. Reich, P.B., Sendall, K.M., Stefanski, A., Wei, X., Rich, R.L., and Montgomery, R.A. (2016). Boreal and temperate trees show strong acclimation of respiration to warming. *Nature* 531, 633–636.
146. Lenhart, K., Weber, B., Elbert, W., Steinkamp, J., Clough, T., Crutzen, P., Pöschl, U., and Keppler, F. (2015). Nitrous oxide and methane emissions from cryptogamic covers. *Global Change Biology* 21, 3889–3900.
147. Skinner, C.B., Poulsen, C.J., and Mankin, J.S. (2018). Amplification of heat extremes by plant CO<sub>2</sub> physiological forcing. *Nature communications* 9, 1–11.

148. Björk, M., Short, F., Mcleod, E., and Beer, S. (2008). Managing seagrasses for resilience to climate change (Iucn).
149. Mcleod, E., Chmura, G.L., Bouillon, S., Salm, R., Björk, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H., and Silliman, B.R. (2011). A blueprint for blue carbon: toward an improved understanding of the role of vegetated coastal habitats in sequestering CO<sub>2</sub>. *Frontiers in Ecology and the Environment* 9, 552–560.
150. Duarte, C.M., Losada, I.J., Hendriks, I.E., Mazarrasa, I., and Marbà, N. (2013). The role of coastal plant communities for climate change mitigation and adaptation. *Nature Climate Change* 3, 961–968.
151. Alongi, D.M. (2015). The impact of climate change on mangrove forests. *Current Climate Change Reports* 1, 30–39.
152. Macreadie, P.I., Nielsen, D.A., Kelleway, J.J., Atwood, T.B., Seymour, J.R., Petrou, K., Connolly, R.M., Thomson, A.C., Trevathan-Tackett, S.M., and Ralph, P.J. (2017). Can we manage coastal ecosystems to sequester more blue carbon? *Frontiers in Ecology and the Environment* 15, 206–213.
153. Boscolo-Galazzo, F., Crichton, K.A., Barker, S., and Pearson, P.N. (2018). Temperature dependency of metabolic rates in the upper ocean: A positive feedback to global climate change? *Global and Planetary Change* 170, 201–212.
154. Park, J.-Y., Kug, J.-S., Bader, J., Rolph, R., and Kwon, M. (2015). Amplified Arctic warming by phytoplankton under greenhouse warming. *Proceedings of the National Academy of Sciences* 112, 5921–5926.
155. Orr, J.C., Fabry, V.J., Aumont, O., Bopp, L., Doney, S.C., Feely, R.A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., et al. (2005). Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature* 437, 681–686.
156. Passow, U., and Carlson, C.A. (2012). The biological pump in a high CO<sub>2</sub> world. *Marine Ecology Progress Series* 470, 249–271.
157. Pörtner, H.-O., Karl, D.M., Boyd, P.W., Cheung, W., Lluch-Cota, S.E., Nojiri, Y., Schmidt, D.N., Zavalov, P.O., Alheit, J., Aristegui, J., et al. (2014). Ocean systems. In *Climate change 2014: impacts, adaptation, and vulnerability. Part A: global and sectoral aspects. contribution of working group II to the fifth assessment report of the intergovernmental panel on climate change* (Cambridge University Press), pp. 411–484.
158. Boyd, P.W. (2015). Toward quantifying the response of the oceans' biological pump to climate change. *Frontiers in Marine Science* 2, 77.
159. Sully, S., Burkepille, D.E., Donovan, M., Hodgson, G., and Van Woesik, R. (2019). A global analysis of coral bleaching over the past two decades. *Nature communications* 10, 1–5.
160. Ayers, G.P., and Cainey, J.M. (2008). The CLAW hypothesis: a review of the major developments. *Environmental Chemistry* 4, 366–374.

161. Quinn, P.K., and Bates, T.S. (2011). The case against climate regulation via oceanic phytoplankton sulphur emissions. *Nature* *480*, 51–56.
162. Six, K.D., Kloster, S., Ilyina, T., Archer, S.D., Zhang, K., and Maier-Reimer, E. (2013). Global warming amplified by reduced sulphur fluxes as a result of ocean acidification. *Nature Climate Change* *3*, 975–978.
163. Wang, S., Maltrud, M., Elliott, S., Cameron-Smith, P., and Jonko, A. (2018). Influence of dimethyl sulfide on the carbon cycle and biological production. *Biogeochemistry* *138*, 49–68.
164. Breider, F., Yoshikawa, C., Makabe, A., Toyoda, S., Wakita, M., Matsui, Y., Kawagucci, S., Fujiki, T., Harada, N., and Yoshida, N. (2019). Response of N<sub>2</sub>O production rate to ocean acidification in the western North Pacific. *Nature Climate Change* *9*, 954–958.
165. Beman, J.M., Chow, C.-E., King, A.L., Feng, Y., Fuhrman, J.A., Andersson, A., Bates, N.R., Popp, B.N., and Hutchins, D.A. (2011). Global declines in oceanic nitrification rates as a consequence of ocean acidification. *Proceedings of the National Academy of Sciences* *108*, 208–213.
166. Collins, W., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C., Joshi, M., Liddicoat, S., et al. (2011). Development and evaluation of an Earth-System model—HadGEM2. *Geoscientific Model Development* *4*, 1051–1075.
167. Knutson, T.R., McBride, J.L., Chan, J., Emanuel, K., Holland, G., Landsea, C., Held, I., Kossin, J.P., Srivastava, A., and Sugi, M. (2010). Tropical cyclones and climate change. *Nature geoscience* *3*, 157–163.
168. Hirabayashi, Y., Mahendran, R., Koirala, S., Konoshima, L., Yamazaki, D., Watanabe, S., Kim, H., and Kanae, S. (2013). Global flood risk under climate change. *Nature Climate Change* *3*, 816–821.
169. Xu, C., Kohler, T.A., Lenton, T.M., Svenning, J.-C., and Scheffer, M. (2020). Future of the human climate niche. *Proceedings of the National Academy of Sciences* *117*, 11350–11355.
170. Kulp, S.A., and Strauss, B.H. (2019). New elevation data triple estimates of global vulnerability to sea-level rise and coastal flooding. *Nature communications* *10*, 1–12.
171. Bronen, R., and Chapin, F.S. (2013). Adaptive governance and institutional strategies for climate-induced community relocations in Alaska. *Proceedings of the National Academy of Sciences* *110*, 9320–9325.
172. Nearing, M., Pruski, F., and O’neal, M. (2004). Expected climate change impacts on soil erosion rates: a review. *Journal of soil and water conservation* *59*, 43–50.
173. Obradovich, N., and Rahwan, I. (2019). Risk of a feedback loop between climatic warming and human mobility. *Journal of the Royal Society Interface* *16*, 20190058.
174. Ho, J. (2010). The implications of Arctic sea ice decline on shipping. *Marine Policy* *34*, 713–715.
175. Humpert, M., and Raspotnik, A. (2012). The future of Arctic shipping. *Port Technology International* *55*, 10–11.

176. Eguíluz, V.M., Fernández-Gracia, J., Irigoien, X., and Duarte, C.M. (2016). A quantitative assessment of Arctic shipping in 2010–2014. *Scientific reports* 6, 1–6.
177. Isaac, M., and Van Vuuren, D.P. (2009). Modeling global residential sector energy demand for heating and air conditioning in the context of climate change. *Energy policy* 37, 507–521.
178. Müller, C., Bondeau, A., Popp, A., Waha, K., and Fader, M. (2010). Climate change impacts on agricultural yields.
179. Bajželj, B., and Richards, K.S. (2014). The positive feedback loop between the impacts of climate change and agricultural expansion and relocation. *Land* 3, 898–916.
180. Thornton, P.E., Calvin, K., Jones, A.D., Di Vittorio, A.V., Bond-Lamberty, B., Chini, L., Shi, X., Mao, J., Collins, W.D., Edmonds, J., et al. (2017). Biospheric feedback effects in a synchronously coupled model of human and Earth systems. *Nature Climate Change* 7, 496–500.
181. Burke, L., Reytar, K., Spalding, M., and Perry, A. (2011). Reefs at risk revisited (World Resources Institute).
182. Frieler, K., Meinshausen, M., Golly, A., Mengel, M., Lebek, K., Donner, S., and Hoegh-Guldberg, O. (2013). Limiting global warming to 2 C is unlikely to save most coral reefs. *Nature Climate Change* 3, 165–170.
183. Hoegh-Guldberg, O., Poloczanska, E.S., Skirving, W., and Dove, S. (2017). Coral reef ecosystems under climate change and ocean acidification. *Frontiers in Marine Science* 4, 158.
184. Misra, A.K. (2014). Climate change and challenges of water and food security. *International Journal of Sustainable Built Environment* 3, 153–165.
185. Koutroulis, A., Papadimitriou, L., Grillakis, M., Tsanis, I., Warren, R., and Betts, R. (2019). Global water availability under high-end climate change: A vulnerability based assessment. *Global and Planetary Change* 175, 52–63.
186. Shahzad, M.W., Burhan, M., and Ng, K.C. (2019). A standard primary energy approach for comparing desalination processes. *npj Clean Water* 2, 1–7.
187. Lykkebo Petersen, K., Heck, N., G Reguero, B., Potts, D., Hovagimian, A., and Paytan, A. (2019). Biological and physical effects of brine discharge from the Carlsbad desalination plant and implications for future desalination plant constructions. *Water* 11, 208.
188. Howard, P., and Livermore, M.A. (2019). Sociopolitical Feedbacks and Climate Change. *Harv. Envtl. L. Rev.* 43, 119.
189. Van Vuuren, D.P., Bayer, L.B., Chuwah, C., Ganzeveld, L., Hazeleger, W., van den Hurk, B., Van Noije, T., O’Neill, B., and Strengers, B.J. (2012). A comprehensive view on climate change: coupling of earth system and integrated assessment models. *Environmental Research Letters* 7, 024012.
190. O’Neill, S., and Nicholson-Cole, S. (2009). “Fear won’t do it” promoting positive engagement with climate change through visual and iconic representations. *Science Communication* 30, 355–379.

191. Moore, F.C., Obradovich, N., Lehner, F., and Baylis, P. (2019). Rapidly declining remarkability of temperature anomalies may obscure public perception of climate change. *Proceedings of the National Academy of Sciences* *116*, 4905–4910.
192. Solaun, K., and Cerdá, E. (2019). Climate change impacts on renewable energy generation. A review of quantitative projections. *Renewable and sustainable energy Reviews* *116*, 109415.
193. Harder, A. (2020). How climate change feeds off itself and gets even worse. *Axios*. <https://www.axios.com/climate-change-feedback-loops-e6cfd8d6-56fe-449b-a856-e8f97717c44c.html>.
194. Woodard, D.L., Davis, S.J., and Randerson, J.T. (2019). Economic carbon cycle feedbacks may offset additional warming from natural feedbacks. *Proceedings of the National Academy of Sciences* *116*, 759–764.
195. Hanson, M. (2020). War is an ecological catastrophe. *Big Think*. <https://bigthink.com/politics-current-affairs/the-environmental-costs-of-war>.
196. Bates, J. (2007). Some considerations of the concept of climate feedback. *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography* *133*, 545–560.
197. Stocker, T.F., Clarke, G.K., Le Treut, H., Lindzen, R.S., Meleshko, V.P., Mugara, R.K., Palmer, T.N., Pierrehumbert, R.T., Sellers, P.J., Trenberth, K.E., et al. (2001). Physical climate processes and feedbacks. In *IPCC, 2001: Climate change 2001: The scientific basis. Contribution of working group I to the third assessment report of the intergovernmental panel on climate change* (Cambridge University Press), pp. 417–470.
198. Wassmann, P., and Lenton, T.M. (2012). Arctic tipping points in an earth system perspective. *Ambio* *41*, 1–9.
199. Kopp, R.E., Shwom, R.L., Wagner, G., and Yuan, J. (2016). Tipping elements and climate–economic shocks: Pathways toward integrated assessment. *Earth’s Future* *4*, 346–372.
200. Wang, S., and Hausfather, Z. (2020). ESD Reviews: mechanisms, evidence, and impacts of climate tipping elements. *Earth System Dynamics Discussions*, 1–93.
201. Rogelj, J., Shindell, D., Jiang, K., Fifita, S., Forster, P., Ginzburg, V., Handa, C., Kheshgi, H., Kobayashi, S., Kriegler, E., et al. (2018). Mitigation pathways compatible with 1.5°C in the context of sustainable development. In *Special Report on the impacts of global warming of 1.5 °C* (Intergovernmental Panel on Climate Change).
202. Rogelj, J., Forster, P.M., Kriegler, E., Smith, C.J., and Séférian, R. (2019). Estimating and tracking the remaining carbon budget for stringent climate targets. *Nature* *571*, 335–342.
203. Forster, P., Rosen, D., Lamboll, R., and Rogelj, J. (2022). Guest post: What the tiny remaining 1.5C carbon budget means for climate policy. *Carbon Brief*. <https://www.carbonbrief.org/guest-post-what-the-tiny-remaining-1-5c-carbon-budget-means-for-climate-policy/>.

204. Friedlingstein, P., O'Sullivan, M., Jones, M.W., Andrew, R.M., Gregor, L., Hauck, J., Le Quéré, C., Lujckx, I.T., Olsen, A., Peters, G.P., et al. (2022). Global carbon budget 2022. *Earth System Science Data* 14, 4811–4900.
205. Caldwell, P.M., Zelinka, M.D., Taylor, K.E., and Marvel, K. (2016). Quantifying the sources of intermodel spread in equilibrium climate sensitivity. *Journal of Climate* 29, 513–524.
206. Colman, R., and Hanson, L. (2017). On the relative strength of radiative feedbacks under climate variability and change. *Climate Dynamics* 49, 2115–2129.
207. Vial, J., Dufresne, J.-L., and Bony, S. (2013). On the interpretation of inter-model spread in CMIP5 climate sensitivity estimates. *Climate Dynamics* 41, 3339–3362.
208. Dessler, A. (2013). Observations of climate feedbacks over 2000–10 and comparisons to climate models. *Journal of Climate* 26, 333–342.
209. Manabe, S., and Wetherald, R.T. (1967). Thermal equilibrium of the atmosphere with a given distribution of relative humidity. *Journal of the Atmospheric Sciences* 24, 241–259.
210. Held, I.M., and Soden, B.J. (2000). Water vapor feedback and global warming. *Annual review of energy and the environment* 25, 441–475.
211. Holland, M.M., and Bitz, C.M. (2003). Polar amplification of climate change in coupled models. *Climate Dynamics* 21, 221–232.
212. Rind, D., Healy, R., Parkinson, C., and Martinson, D. (1995). The role of sea ice in 2×CO<sub>2</sub> climate model sensitivity. Part I: The total influence of sea ice thickness and extent. *Journal of Climate* 8, 449–463.
213. Robinson, A.J., and Goelzer, H. (2014). The importance of insolation changes for paleo ice sheet modeling. *Cryosphere* 8, 1419–1428.
214. Tedesco, M., Doherty, S., Fettweis, X., Alexander, P., Jeyaratnam, J., and Stroeve, J. (2016). The darkening of the Greenland ice sheet: trends, drivers, and projections (1981–2100).
215. Tedstone, A.J., Bamber, J.L., Cook, J.M., Williamson, C.J., Fettweis, X., Hodson, A.J., and Tranter, M. (2017). Dark ice dynamics of the south-west Greenland Ice Sheet. *Cryosphere* 11, 2491–2506.
216. Ridley, J., Gregory, J.M., Huybrechts, P., and Lowe, J. (2010). Thresholds for irreversible decline of the Greenland ice sheet. *Climate Dynamics* 35, 1049–1057.
217. Köhler, P., Bintanja, R., Fischer, H., Joos, F., Knutti, R., Lohmann, G., and Masson-Delmotte, V. (2010). What caused Earth's temperature variations during the last 800,000 years? Data-based evidence on radiative forcing and constraints on climate sensitivity. *Quaternary Science Reviews* 29, 129–145.
218. Budyko, M.I. (1969). The effect of solar radiation variations on the climate of the Earth. *tellus* 21, 611–619.

219. Sellers, W.D. (1969). A global climatic model based on the energy balance of the earth-atmosphere system. *Journal of Applied Meteorology and Climatology* 8, 392–400.
220. Schneider, S.H., and Dickinson, R.E. (1974). Climate modeling. *Reviews of Geophysics* 12, 447–493.
221. Robock, A. (1983). Ice and snow feedbacks and the latitudinal and seasonal distribution of climate sensitivity. *Journal of the Atmospheric Sciences* 40, 986–997.
222. Robock, A. (1985). An updated climate feedback diagram. *Bulletin of the American Meteorological Society* 66, 786–787.
223. Cess, R. o, Potter, G., Zhang, M.-H., Blanchet, J.-P., Chalita, S., Colman, R., Dazlich, D., Del Genio, A., Dymnikov, V., Galin, V., et al. (1991). Interpretation of snow-climate feedback as produced by 17 general circulation models. *Science* 253, 888–892.
224. Randall, D. o, Cess, R., Blanchet, J., Chalita, S., Colman, R., Dazlich, D., Del Genio, A., Keup, E., Lacis, A., Le Treut, H., et al. (1994). Analysis of snow feedbacks in 14 general circulation models. *Journal of Geophysical Research: Atmospheres* 99, 20757–20771.
225. Hall, A. (2004). The role of surface albedo feedback in climate. *Journal of Climate* 17, 1550–1568.
226. Winton, M. (2006). Surface albedo feedback estimates for the AR4 climate models. *Journal of Climate* 19, 359–365.
227. Flanner, M.G., Shell, K.M., Barlage, M., Perovich, D.K., and Tschudi, M. (2011). Radiative forcing and albedo feedback from the Northern Hemisphere cryosphere between 1979 and 2008. *Nature Geoscience* 4, 151–155.
228. Fletcher, C.G., Zhao, H., Kushner, P.J., and Fernandes, R. (2012). Using models and satellite observations to evaluate the strength of snow albedo feedback. *Journal of Geophysical Research: Atmospheres* 117.
229. Robock, A. (1980). The seasonal cycle of snow cover, sea ice and surface albedo. *Monthly Weather Review* 108, 267–285.
230. Prospero, J.M., and Lamb, P.J. (2003). African droughts and dust transport to the Caribbean: Climate change implications. *Science* 302, 1024–1027.
231. Moulin, C., and Chiapello, I. (2006). Impact of human-induced desertification on the intensification of Sahel dust emission and export over the last decades. *Geophysical Research Letters* 33.
232. Gettelman, A., Lin, L., Medeiros, B., and Olson, J. (2016). Climate feedback variance and the interaction of aerosol forcing and feedbacks. *Journal of Climate* 29, 6659–6675.
233. De Lavergne, C., Palter, J.B., Galbraith, E.D., Bernardello, R., and Marinov, I. (2014). Cessation of deep convection in the open Southern Ocean under anthropogenic climate change. *Nature Climate Change* 4, 278–282.

234. Breitburg, D., Levin, L.A., Oschlies, A., Grégoire, M., Chavez, F.P., Conley, D.J., Garçon, V., Gilbert, D., Gutiérrez, D., Isensee, K., et al. (2018). Declining oxygen in the global ocean and coastal waters. *Science* 359.
235. DeVries, T., Holzer, M., and Primeau, F. (2017). Recent increase in oceanic carbon uptake driven by weaker upper-ocean overturning. *Nature* 542, 215–218.
236. Bindoff, N.L., Cheung, W.W.L., Kairo, J.G., Arístegui, J., Gunder, V.A., Hallberg, R., Hilmi, N., Jiao, N., Karim, M. saiful, Levin, L., et al. (2019). Changing Ocean, Marine Ecosystems, and Dependent Communities. In IPCC special report on the ocean and cryosphere in a changing climate, H.-O. Pörtner, D. C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, M. Nicolai, A. Okem, J. Petzold, et al., eds.
237. Balaguru, K., Foltz, G.R., Leung, L.R., and Emanuel, K.A. (2016). Global warming-induced upper-ocean freshening and the intensification of super typhoons. *Nature communications* 7, 1–8.
238. Keeling, R.F., Körtzinger, A., and Gruber, N. (2009). Ocean deoxygenation in a warming world.
239. Fu, W., Randerson, J.T., and Moore, J.K. (2016). Climate change impacts on net primary production (NPP) and export production (EP) regulated by increasing stratification and phytoplankton community structure in the CMIP5 models. *Biogeosciences* 13, 5151–5170.
240. Boyce, D.G., Lewis, M.R., and Worm, B. (2010). Global phytoplankton decline over the past century. *Nature* 466, 591–596.
241. Vellinga, M., and Wood, R.A. (2002). Global climatic impacts of a collapse of the Atlantic thermohaline circulation. *Climatic change* 54, 251–267.
242. Zhang, R., and Delworth, T.L. (2005). Simulated tropical response to a substantial weakening of the Atlantic thermohaline circulation. *Journal of climate* 18, 1853–1860.
243. Stouffer, R.J., Yin, J., Gregory, J., Dixon, K., Spelman, M., Hurlin, W., Weaver, A., Eby, M., Flato, G., Hasumi, H., et al. (2006). Investigating the causes of the response of the thermohaline circulation to past and future climate changes. *Journal of Climate* 19, 1365–1387.
244. Stone, P.H., and Carlson, J.H. (1979). Atmospheric lapse rate regimes and their parameterization. *Journal of the Atmospheric Sciences* 36, 415–423.
245. Xu, K.-M., and Emanuel, K.A. (1989). Is the tropical atmosphere conditionally unstable? *Monthly Weather Review* 117, 1471–1479.
246. Oerlemans, J. (1981). Some basic experiments with a vertically-integrated ice sheet model. *Tellus* 33, 1–11.
247. Hakuba, M.Z., Folini, D., Wild, M., and Schär, C. (2012). Impact of Greenland’s topographic height on precipitation and snow accumulation in idealized simulations. *Journal of Geophysical Research: Atmospheres* 117.



248. Ridley, J.K., Huybrechts, P., Gregory, J. u, and Lowe, J. (2005). Elimination of the Greenland ice sheet in a high CO2 climate. *Journal of Climate* 18, 3409–3427.
249. Gregoire, L.J., Otto-Bliesner, B., Valdes, P.J., and Ivanovic, R. (2016). Abrupt Bølling warming and ice saddle collapse contributions to the Meltwater Pulse 1a rapid sea level rise. *Geophysical research letters* 43, 9130–9137.
250. Van der Veen, C. (1998). Fracture mechanics approach to penetration of surface crevasses on glaciers. *Cold Regions Science and Technology* 27, 31–47.
251. DeConto, R.M., and Pollard, D. (2016). Contribution of Antarctica to past and future sea-level rise. *Nature* 531, 591–597.
252. Phipps, S.J., Fogwill, C.J., and Turney, C.S. (2016). Impacts of marine instability across the East Antarctic Ice Sheet on Southern Ocean dynamics. *The Cryosphere* 10, 2317–2328.
253. Notz, D., and Marotzke, J. (2012). Observations reveal external driver for Arctic sea-ice retreat. *Geophysical Research Letters* 39.
254. Hezel, P., Zhang, X., Bitz, C., Kelly, B., and Massonnet, F. (2012). Projected decline in spring snow depth on Arctic sea ice caused by progressively later autumn open ocean freeze-up this century. *Geophysical Research Letters* 39.
255. Tietsche, S., Notz, D., Jungclaus, J., and Marotzke, J. (2011). Recovery mechanisms of Arctic summer sea ice. *Geophysical Research Letters* 38.
256. Taylor, K.E., Stouffer, R.J., and Meehl, G.A. (2012). An overview of CMIP5 and the experiment design. *Bulletin of the American Meteorological Society* 93, 485–498.
257. Kravitz, B., Robock, A., Forster, P.M., Haywood, J.M., Lawrence, M.G., and Schmidt, H. (2013). An overview of the Geoengineering Model Intercomparison Project (GeoMIP). *Journal of Geophysical Research: Atmospheres* 118, 13,103-13,107. <https://doi.org/10.1002/2013JD020569>.
258. Cionni, I., Eyring, V., Lamarque, J.-F., Randel, W., Stevenson, D., Wu, F., Bodeker, G., Shepherd, T., Shindell, D., and Waugh, D. (2011). Ozone database in support of CMIP5 simulations: results and corresponding radiative forcing. *Atmospheric Chemistry and Physics* 11, 11267–11292.
259. Eyring, V., Arblaster, J.M., Cionni, I., Sedláček, J., Perlwitz, J., Young, P.J., Bekki, S., Bergmann, D., Cameron-Smith, P., Collins, W.J., et al. (2013). Long-term ozone changes and associated climate impacts in CMIP5 simulations. *Journal of Geophysical Research: Atmospheres* 118, 5029–5060.
260. Jones, Cd., Hughes, J., Bellouin, N., Hardiman, S., Jones, G., Knight, J., Liddicoat, S., O’connor, F., Andres, R.J., Bell, C., et al. (2011). The HadGEM2-ES implementation of CMIP5 centennial simulations. *Geoscientific Model Development* 4, 543–570.
261. Stevenson, D., Young, P., Naik, V., Lamarque, J.-F., Shindell, D.T., Voulgarakis, A., Skeie, R.B., Dalsoren, S.B., Myhre, G., Berntsen, T.K., et al. (2013). Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP). *Atmospheric Chemistry and Physics* 13, 3063–3085.

262. Dietmüller, S., Ponater, M., and Sausen, R. (2014). Interactive ozone induces a negative feedback in CO<sub>2</sub>-driven climate change simulations. *Journal of Geophysical Research: Atmospheres* *119*, 1796–1805.
263. Prather, M.J. (1996). Time scales in atmospheric chemistry: Theory, GWPs for CH<sub>4</sub> and CO, and runaway growth. *Geophysical Research Letters* *23*, 2597–2600.
264. Stevenson, D., Dentener, F., Schultz, M., Ellingsen, K., Van Noije, T., Wild, O., Zeng, G., Amann, M., Atherton, C., Bell, N., et al. (2006). Multimodel ensemble simulations of present-day and near-future tropospheric ozone. *Journal of Geophysical Research: Atmospheres* *111*.
265. Fowler, D., Amann, M., Anderson, R., Ashmore, M., Cox, P., Depledge, M., Derwent, D., Grennfelt, P., Hewitt, N., Hov, O., et al. (2008). Ground-level ozone in the 21st century: future trends, impacts and policy implications.
266. Garrels, R.M., Mackenzie, F.T., and Hunt, C. (1973). Chemical cycles and the global environment: assessing human influences.
267. Olefeldt, D., Euskirchen, E.S., Harden, J., Kane, E., McGuire, A.D., Waldrop, M.P., and Turetsky, M.R. (2017). A decade of boreal rich fen greenhouse gas fluxes in response to natural and experimental water table variability. *Global change biology* *23*, 2428–2440.
268. Collins, M., Knutti, R., Arblaster, J., Dufresne, J.-L., Fichet, T., Friedlingstein, P., Gao, X., Gutowski, W.J., Johns, T., Krinner, G., et al. (2013). Long-term climate change: projections, commitments and irreversibility. In *Climate Change 2013-The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (Cambridge University Press), pp. 1029–1136.
269. Mitsch, W.J., Nahlik, A., Wolski, P., Bernal, B., Zhang, L., and Ramberg, L. (2010). Tropical wetlands: seasonal hydrologic pulsing, carbon sequestration, and methane emissions. *Wetlands ecology and management* *18*, 573–586.
270. Glaser, P.H., Siegel, D.I., Chanton, J.P., Reeve, A.S., Rosenberry, D.O., Corbett, J.E., Dasgupta, S., and Levy, Z. (2016). Climatic drivers for multidecadal shifts in solute transport and methane production zones within a large peat basin. *Global biogeochemical cycles* *30*, 1578–1598.
271. Serreze, M.C., and Barry, R.G. (2011). Processes and impacts of Arctic amplification: A research synthesis. *Global and planetary change* *77*, 85–96.
272. Chapin, F.S., Sturm, M., Serreze, M.C., McFadden, J.P., Key, J., Lloyd, A.H., McGuire, A., Rupp, T.S., Lynch, A.H., Schimel, J.P., et al. (2005). Role of land-surface changes in Arctic summer warming. *science* *310*, 657–660.
273. Thompson, C., Beringer, J., Chapin III, F.S., and McGuire, A.D. (2004). Structural complexity and land-surface energy exchange along a gradient from arctic tundra to boreal forest. *Journal of Vegetation Science* *15*, 397–406.

274. Beringer, J., Chapin III, F.S., Thompson, C.C., and McGuire, A.D. (2005). Surface energy exchanges along a tundra-forest transition and feedbacks to climate. *Agricultural and Forest Meteorology* *131*, 143–161.
275. Pearson, R.G., Phillips, S.J., Loranty, M.M., Beck, P.S., Damoulas, T., Knight, S.J., and Goetz, S.J. (2013). Shifts in Arctic vegetation and associated feedbacks under climate change. *Nature climate change* *3*, 673–677.
276. Déry, S.J., and Brown, R.D. (2007). Recent Northern Hemisphere snow cover extent trends and implications for the snow-albedo feedback. *Geophysical Research Letters* *34*.
277. Matthes, H., Rinke, A., Miller, P.A., Kuhry, P., Dethloff, K., and Wolf, A. (2012). Sensitivity of high-resolution Arctic regional climate model projections to different implementations of land surface processes. *Climatic Change* *111*, 197–214.
278. Kurz, W., Apps, M., Webb, T., and McNamee, P. (1992). Carbon budget of the Canadian forest sector. Phase I. Forestry Canada. Northern Forestry Centre, Edmonton (Inf. Rep. NOR-X-326).
279. Kurz, W.A., and Apps, M.J. (1999). A 70-year retrospective analysis of carbon fluxes in the Canadian forest sector. *Ecological applications* *9*, 526–547.
280. Kurz, W., and Apps, M. (2006). Developing Canada's national forest carbon monitoring, accounting and reporting system to meet the reporting requirements of the Kyoto Protocol. *Mitigation and Adaptation Strategies for Global Change* *11*, 33–43.
281. Kulmala, M., Suni, T., Lehtinen, K.E.J., Dal Maso, M., Boy, M., Reissell, A., Rannik, Ü., Aalto, P., Keronen, P., Hakola, H., et al. (2004). A new feedback mechanism linking forests, aerosols, and climate. *Atmospheric Chemistry and Physics* *4*, 557–562. 10.5194/acp-4-557-2004.
282. Guenther, A., Hewitt, C.N., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W., et al. (1995). A global model of natural volatile organic compound emissions. *Journal of Geophysical Research: Atmospheres* *100*, 8873–8892.
283. Arneth, A., Niinemets, Ü., Pressley, S., Bäck, J., Hari, P., Karl, T., Noe, S., Prentice, I.C., Serça, D., Hickler, T., et al. (2007). Process-based estimates of terrestrial ecosystem isoprene emissions: incorporating the effects of a direct CO<sub>2</sub>-isoprene interaction. *Atmospheric Chemistry and Physics* *7*, 31–53. 10.5194/acp-7-31-2007.
284. Arneth, A., Miller, P.A., Scholze, M., Hickler, T., Schurgers, G., Smith, B., and Prentice, I.C. (2007). CO<sub>2</sub> inhibition of global terrestrial isoprene emissions: Potential implications for atmospheric chemistry. *Geophysical Research Letters* *34*.
285. Palmer, P.I., Abbot, D.S., Fu, T.-M., Jacob, D.J., Chance, K., Kurosu, T.P., Guenther, A., Wiedinmyer, C., Stanton, J.C., Pilling, M.J., et al. (2006). Quantifying the seasonal and interannual variability of North American isoprene emissions using satellite observations of the formaldehyde column. *Journal of Geophysical Research: Atmospheres* *111*.
286. Bellamy, P.H., Loveland, P.J., Bradley, R.I., Lark, R.M., and Kirk, G.J. (2005). Carbon losses from all soils across England and Wales 1978–2003. *Nature* *437*, 245–248.

287. Davidson, E.A., and Janssens, I.A. (2006). Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature* *440*, 165–173.
288. Billings, W. (1987). Carbon balance of Alaskan tundra and taiga ecosystems: past, present and future. *Quaternary Science Reviews* *6*, 165–177.
289. Jenkinson, D.S., Adams, D., and Wild, A. (1991). Model estimates of CO<sub>2</sub> emissions from soil in response to global warming. *Nature* *351*, 304–306.
290. Menon, S., Denman, K.L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P.M., Dickinson, R.E., Hauglustaine, D., Heinze, C., Holland, E., et al. (2007). Couplings between changes in the climate system and biogeochemistry (Lawrence Berkeley National Lab.(LBNL), Berkeley, CA (United States)).
291. Barnard, R., Leadley, P.W., and Hungate, B.A. (2005). Global change, nitrification, and denitrification: a review. *Global biogeochemical cycles* *19*.
292. Li, C., Frohling, S., and Frohling, T.A. (1992). A model of nitrous oxide evolution from soil driven by rainfall events: 2. Model applications. *Journal of Geophysical Research: Atmospheres* *97*, 9777–9783.
293. Smith, K. (1997). The potential for feedback effects induced by global warming on emissions of nitrous oxide by soils. *Global Change Biology* *3*, 327–338.
294. Kesik, M., Brüggemann, N., Forkel, R., Kiese, R., Knoche, R., Li, C., Seufert, G., Simpson, D., and Butterbach-Bahl, K. (2006). Future scenarios of N<sub>2</sub>O and NO emissions from European forest soils. *Journal of Geophysical Research: Biogeosciences* *111*.
295. Xu-Ri, and Prentice, I.C. (2008). Terrestrial nitrogen cycle simulation with a dynamic global vegetation model. *Global Change Biology* *14*, 1745–1764.
296. Khalil, M.A.K., and Rasmussen, R. (1989). Climate-induced feedbacks for the global cycles of methane and nitrous oxide. *Tellus B: Chemical and Physical Meteorology* *41*, 554–559.
297. Seneviratne, S.I., Corti, T., Davin, E.L., Hirschi, M., Jaeger, E.B., Lehner, I., Orlowsky, B., and Teuling, A.J. (2010). Investigating soil moisture–climate interactions in a changing climate: A review. *Earth-Science Reviews* *99*, 125–161.
298. Farquhar, G.D., von Caemmerer, S. von, and Berry, J.A. (1980). A biochemical model of photosynthetic CO<sub>2</sub> assimilation in leaves of C<sub>3</sub> species. *Planta* *149*, 78–90.
299. Woodward, F.I. (1987). Stomatal numbers are sensitive to increases in CO<sub>2</sub> from pre-industrial levels. *Nature* *327*, 617–618.
300. Willeit, M., Ganopolski, A., and Feulner, G. (2014). Asymmetry and uncertainties in biogeophysical climate–vegetation feedback over a range of CO<sub>2</sub> forcings. *Biogeosciences* *11*, 17–32.

301. Breshears, D.D., Adams, H.D., Eamus, D., McDowell, N., Law, D.J., Will, R.E., Williams, A.P., and Zou, C.B. (2013). The critical amplifying role of increasing atmospheric moisture demand on tree mortality and associated regional die-off. *Frontiers in plant science* 4, 266.
302. Teskey, R., Wertin, T., Bauweraerts, I., Ameye, M., McGuire, M.A., and Steppe, K. (2015). Responses of tree species to heat waves and extreme heat events. *Plant, cell & environment* 38, 1699–1712.
303. Matusick, G., Ruthrof, K.X., Brouwers, N.C., Dell, B., and Hardy, G.S.J. (2013). Sudden forest canopy collapse corresponding with extreme drought and heat in a mediterranean-type eucalypt forest in southwestern Australia. *European Journal of Forest Research* 132, 497–510.
304. Villegas, J.C., Law, D.J., Stark, S.C., Minor, D.M., Breshears, D.D., Saleska, S.R., Swann, A.L., Garcia, E.S., Bella, E.M., Morton, J.M., et al. (2017). Prototype campaign assessment of disturbance-induced tree loss effects on surface properties for atmospheric modeling. *Ecosphere* 8, e01698.
305. Pendleton, L., Donato, D.C., Murray, B.C., Crooks, S., Jenkins, W.A., Sifleet, S., Craft, C., Fourqurean, J.W., Kauffman, J.B., Marbà, N., et al. (2012). Estimating global “blue carbon” emissions from conversion and degradation of vegetated coastal ecosystems. *PloS one* 7, e43542.
306. Volk, T., and Hoffert, M.I. (1985). Ocean carbon pumps: Analysis of relative strengths and efficiencies in ocean-driven atmospheric CO<sub>2</sub> changes. *The carbon cycle and atmospheric CO<sub>2</sub>: natural variations Archean to present* 32, 99–110.
307. Boyd, P., and Stevens, C. (2002). Modelling particle transformations and the downward organic carbon flux in the NE Atlantic Ocean. *Progress in Oceanography* 52, 1–29.
308. Huang, L., Krigsvoll, G., Johansen, F., Liu, Y., and Zhang, X. (2018). Carbon emission of global construction sector. *Renewable and Sustainable Energy Reviews* 81, 1906–1916.
309. Geisler, C., and Currens, B. (2017). Impediments to inland resettlement under conditions of accelerated sea level rise. *Land Use Policy* 66, 322–330.
310. Hoffmann, R., Dimitrova, A., Muttarak, R., Cuaresma, J.C., and Peisker, J. (2020). A meta-analysis of country-level studies on environmental change and migration. *Nature Climate Change* 10, 904–912.
311. Liang, S., Yang, X., Qi, J., Wang, Y., Xie, W., Muttarak, R., and Guan, D. (2020). CO<sub>2</sub> Emissions Embodied in International Migration from 1995 to 2015. *Environmental Science & Technology* 54, 12530–12538.
312. Stehfest, E., Bouwman, L., Van Vuuren, D.P., Den Elzen, M.G., Eickhout, B., and Kabat, P. (2009). Climate benefits of changing diet. *Climatic change* 95, 83–102.
313. Carlsson-Kanyama, A., and González, A.D. (2009). Potential contributions of food consumption patterns to climate change. *The American journal of clinical nutrition* 89, 1704S-1709S.
314. Norgaard, K.M. (2011). *Living in denial: Climate change, emotions, and everyday life* (MIT Press).

315. Horowitz, J.K. (2009). The income–temperature relationship in a cross-section of countries and its implications for predicting the effects of global warming. *Environmental and Resource economics* 44, 475–493.
316. Weitzman, M.L. (2009). On modeling and interpreting the economics of catastrophic climate change. *The Review of Economics and Statistics* 91, 1–19.